

HERBICIDE
PRE-PLANT, PRE-EMERGENCEGROUP
2GROUP
5MULTI
MODE
OF ACTION

CANOPY PRO APPLICATION GUIDELINES

Canopy™ PRO**HERBICIDE**

Canopy™ PRO pre-emergence herbicide delivers broad-spectrum and residual activity to help maximize early-season control of tough weeds.

WHY USE CANOPY PRO HERBICIDE?

- **Broad spectrum** weed control performance
- Control resistant **Canada fleabane**
- **Optimize soybean yield** by removing weeds early

WEEDS CONTROLLED**Field corn (25 g/ac)**

Broadleaf weeds

- Buckwheat, Wild
- Canada Fleabane⁴
- Carpetweed
- Chickweed, Common
- Cocklebur¹
- Corn Spurry
- Dandelion
- Hemp-nettle
- Jimsonweed¹
- Lady's-thumb
- Lamb's-quarters, Common
- Mustard, Wild
- Pigweed, Prostrate
- Pigweed, Redroot
- Potato Vine, Wild

• Ragweed, Common

• Shepherd's-purse

• Smartweed, Green

• Sow-thistle, Annual⁵

• Thistle, Russian

• Velvetleaf

• Woodsorrel, Yellow

Annual grasses• Barnyard Grass²• Cheatgrass²• Crabgrass²• Foxtail, Giant²• Foxtail, Green²• Foxtail, Yellow²• Johnsongrass²• Nutsedge, Yellow³• Panicum, Fall²• Witchgrass²

Soybeans

CROPS

Soybeans

RATES AND ACRES TREATED**Rates:**

• 14.4 g/ac + 225 g/ac

Acres treated:

• New 40 ac/case

Water volume:

• 10-20 US gal/ac

PACKAGING**Case:**

• Classic 2 x 288 g
bottles + Metribuzin Mx
2 x 4.5 kg jugs

WHEN TO APPLY**PRE:** Apply after planting but prior to crop emergence**Pre-Plant Burndown:** Apply up to 14 days before planting**RAINFAST**

• 2 hours

HERBICIDE TANK MIXES**Registered:**

2,4-D Ester
Dual II Magnum[®]
Enlist Duo™ or Enlist™ 1[®]

Eragon[®] LQ
Frontier[®] MAX
Glyphosate

[®] Only for use in Enlist E3™ soybeans**Application information**

Formulated as a dry flowable granule to be mixed in water and applied as uniform broadcast spray. Only apply using ground equipment. This product is not registered for aerial application.

Crop rotation

The recropping intervals are dependent on the pH of the soil, as well as the sensitivity of the specific rotational crop. Consult product labels for rotation intervals across a full range of soil pH levels.

Soil pH ≤ to 7.4:

3 months: winter wheat**10 months:** field corn, soybeans, white beans and alfalfa**12 months:** tomatoes

See product label for full list of crop rotation guidelines.

Pre-harvest interval

- The PHI is 60 days.

Mixing Instructions

1. Fill the sprayer tank with ½ to ¾ of the required amount of water
2. Continue agitation throughout the mixing and spraying procedure
3. Add the required amount of Classic™ and allow time for product to fully disperse
4. Add the required amount of Metribuzin Mx
5. If required for a burndown, add liquids or suspension concentrates such as Enlist 1 or Enlist Duo
6. If required for a burndown, add glyphosate and continue agitation
7. Complete filling the sprayer tank

¹ Large-seeded weeds that germinate deep in the soil, such as cocklebur and jimsonweed, may not be fully controlled.

² Partial control

³ Suppression

⁴ Including glyphosate-resistant biotypes. Must be tank mixed with Eragon[®]. For RoundupReady 2 Xtend™ soybeans, tank mix with a low volatile dicamba herbicide. For Enlist E3™ soybeans, tank mix with Enlist Duo™ or Enlist™ 1.

⁵ Must be tank mixed with glyphosate