

For strong pre-emergent residual control of the toughest broadleaf weeds in herbicide-tolerant soybeans.

# **Registered crops**

Soybeans

# Weeds controlled

- · Barnyard grass
- Canada fleabane (including Group 2-, 5and glyphosateresistant biotypes)
- Carpetweed<sup>1</sup>
- Cheatgrass
- Chickweed
- Cleavers (suppression)
- Cocklebur
- Common ragweed (including Group 2-, 5and glyphosateresistant biotypes)
- Crabgrass
- Dandelion
- Eastern black nightshade (including Group 2- and 22-resistant biotypes)

- Fall panicum
- Giant foxtail
- Green and redroot pigweed (including Group 2- and 5-resistant biotypes)
- Green smartweed
- Green and yellow foxtail
- · Hairy nightshade
- Jimsonweed<sup>1</sup>
- Johnson grass (seedling)
- · Lady's-thumb
- Lamb's-quarters (including Group 2- and 5-resistant biotypes)
- Palmer amaranth
- Prickly mallow<sup>2</sup>

- · Prostrate pigweed
- · Russian thistle
- · Shepherd's-purse
- Velvetleaf
- Volunteer canola (suppression including all herbicide-tolerant biotypes)
- Waterhemp (including Group 2-, 5and glyphosateresistant biotypes)
- · Wild buckwheat
- Wild mustard
- · Wild potato vine
- Witchgrass
- Yellow woodsorrel

## **ACTIVE INGREDIENTS**

- > Group 5 (metribuzin)
- > Group 14 (flumioxazin)

## TREATMENT AREA

> One case treats 40 acres

#### APPLICATION TIMING

> Pre-emergent for soybeans

### **TANK-MIX ORDER**

- > Bifecta EZ
- > BlackHawk (if used)
- > Glyphosate

#### PCP#

> 34422

# **Benefits**

- STRONG GLYPHOSATE-RESISTANT fleabane burndown and control when tank-mixed with BlackHawk® herbicide
- RESIDUAL WEED CONTROL for up to 5 weeks
- SUPERIOR RESISTANCE MANAGEMENT with two active ingredients to help manage weeds that are Group 2-, 5- and glyphosate-resistant
- Convenient all-in-one formulation
- Tank-mixable with most burndown and pre-emergent herbicides (i.e. glyphosate and BlackHawk)
- · Flexible pre-plant or pre-emergent application timings
- No surfactant needed
- <sup>1</sup> Pre-emergence only
- <sup>2</sup> Post-emergence only

