

Tips for Providing Safe Structure for Adult Behavioral Health Patients

It is well documented that many behavioral health patients in the emergency department (ED) waiting to be admitted have extended lengths of stay, sometimes days or even weeks due to a lack of available beds either within the facility or at an external facility. In such cases, emergency nurses can ensure safety and structure for this patient population by providing these patients with a daily routine. These infographics have been developed to promote conversation between emergency nurses and the ED team as a starting point to help the ED team create and provide a daily routine for ED behavioral health patients that is safe, structured, and adheres to individual institutional policies.

Suggested daily routine:









Encourage a supportive person to stay with the patient.

Inform that person of what will take place while in the emergency department.



Provide 1:1 observation if the patient is at risk for suicide (The Joint Commission, 2016).



Search the patient's clothing and belongings and remove anything that is potentially harmful.



Monitor visitors as needed. Ensure the patient's communication with visitors is therapeutic.



Monitor the use of personal electronic devices and be prepared to take them away if necessary.



Check the patient's room periodically.
Remove dangerous items such as cans,
glass, sharp objects, plastic bags, harmful
liquids, metal/sharp hair accessories,
belts, string/ribbon/cords, knives/scissors,
lighters/matches, medicines, etc.

ENA Infographics are informational documents developed by ENA members to provide the reader with knowledge on a subject relevant to emergency care and are made available for educational and information purposes only. The information and recommendations contained in this infographic reflect current knowledge at the time of publication, are only current as of its publication date, and are subject to change without notice as advances emerge. Additionally, variations in practice, which take into account the needs of the individual patient or institution and the resources and limitations unique to the health care setting, may warrant approaches, treatments and/or procedures that



differ from the recommendations outlined in this infographic. Therefore, these recommendations should not be interpreted as dictating an exclusive course of management, treatment or care, nor does the use of such recommendations guarantee a particular outcome. Infographics are never intended to replace a practitioner's best nursing judgment based on the clinical circumstances of a particular patient or patient population. ENA does not "approve" or "endorse" any specific methods, practices or sources of information. ENA assumes no liability for any injury and/or damage to persons or property arising out of or related to the use of or reliance on this infographic.