

Enhance Recovery After Surgery: ERAS


Matthew A. Barker, MD, FACOG
Avera Medical Group – Urogynecology

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



1

Matthew A. Barker, MD



- The University of South Dakota Sanford School of Medicine
 - Professor of ObGyn, Internal Medicine, and Neurosciences
- Avera McKennan Hospital & University Health Center
 - Director of Female Pelvic Medicine & Reconstructive Surgery
 - Avera Medical Group - Urogynecology
- Education:
 - BS: Creighton University, Omaha, NE
 - MD: The University of South Dakota Sanford School of Medicine
 - ObGyn Residency: University of Wisconsin – Madison, WI
 - Urogynecology Fellowship: Good Samaritan Hospital Cincinnati, OH
- Board Certified
 - Obstetrics and Gynecology
 - Urogynecology and Pelvic Reconstructive Surgery

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



2

Disclosures

- Speaker for Astellas
- Speaker for Axonics

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



3

Objectives

- Learn the benefits of ERAS in perioperative management of surgical patients.
- Understand why surgeons utilize ERAS and how to implement these changes at your institution.

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



4

Questions?

- Does ERAS decrease the risk of complications in surgery? True or False
- Glycemic control is **NOT** important for minimizing complications during the perioperative period for surgical patients? True or False

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



5

"Wait and see"
approach

Team reacts to
postoperative
events



More active
approach


Referred to as
ERAS or "fast
track" protocol

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



6


Impact of Surgery



AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

7
Avera
Medical Group
Urogynecology

Goal of ERAS



AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

8
Avera
Medical Group
Urogynecology

Phases of Care: PREoperative

- Thorough counseling and education
- Optimization of co-morbid conditions
- No mechanical bowel preparation
- No overnight fasting
- Oral carbohydrate loading
- Avoidance of long-acting sedatives
- Acetaminophen, gabapentin, Celecoxib




Scheib et al, JMIG, 2019

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

9
Avera
Medical Group
Urogynecology

Phases of Care: INTRAoperative

- Antimicrobial prophylaxis
- Thrombosis prophylaxis
- Routine antiemetics
- Epidural analgesia
- High O2 concentrations
- Preventing hypothermia
- Avoidance of pelvic drains
- Goal-directed IV fluids/euvolemia
- Wound infiltration with local anesthetic/nerve blocks



Scheib et al, JMIG, 2019

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

10
Avera
Medical Group
Urogynecology

Phases of Care: POSToperative

- Avoidance of nasogastric tubes
- Avoidance of ileus
- Prevention of PONV
- Multimodal analgesia
- Early oral intake
- Nutritional supplements
- Early mobilization, physical therapy
- Thrombosis prophylaxis



Scheib et al, JMIG, 2019

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

11
Avera
Medical Group
Urogynecology

ERAS Works!

- **Length of stay↓:**
 - TAH pre-protocol 5 days → post-protocol 3 days
 - TLH pre-protocol same day discharge 9% → post-protocol 56%
- **Postop pain and narcotic use↓:** 20.8% to 97.4% drop in narcotic use
- **Bowel function:** First defecation 24 hours sooner in ERAS group
- **Ambulation:** 75.6% ambulating within 3 hours of surgery
- **Complications or Readmissions:** No significant difference

Mukhopadhyay et al, BMJ, 2015; Keil et al, Anesth, Analg, 2018; Hansen et al, AJOG, 2007

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

12
Avera
Medical Group
Urogynecology

ERAS Works!

- **Patient satisfaction**

- Positive outcomes for abdominal, laparoscopic, and vaginal procedures
- LaPasse et al evaluated patients at POD 7 and POD 30 → 97% reported satisfaction with their care

- **Economics cost analysis**

- ERAS implementation has been shown to decrease average hospital costs by 18%-21% per patient

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



13



AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



14

Developing your ERAS Protocol

Guidelines for pre- and intra-operative care in gynecologic/oncology surgery: Enhanced Recovery After Surgery (ERAS®) Society recommendations — Part I

Nelson G.^{1,2}, Altman A.D.³, Nick A.⁴, Meyer L.A.⁵, Ramirez P.T.⁶, Ahtari C.⁷, Antrobus J.⁸, Huang J.⁹, Scott M.¹⁰, Wijk L.¹, Acheson N.¹, Ljungqvist O.¹, Dowdy S.C.¹

Guidelines for postoperative care in gynecologic/oncology surgery: Enhanced Recovery After Surgery (ERAS®) Society recommendations — Part II

G. Nelson^{1,2}, A.D. Altman³, A. Nick⁴, L.A. Meyer⁵, P.T. Ramirez⁶, C. Ahtari⁷, J. Antrobus⁸, J. Huang⁹, M. Scott¹⁰, L. Wijk¹, N. Acheson¹, O. Ljungqvist¹, S.C. Dowdy¹

Enhanced Recovery after Surgery in Gynecology: A Review of the Literature



Stacey A. Scheib, MD, May Thomasson, MD, and Juanan L. Kenner, MD
From the Department of Obstetrics and Gynecology, Louisiana State University Health Sciences Center, New Orleans, Louisiana (all authors)

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



15

Our Experience




AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



16

Key Stakeholders



AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



17

Identify a champion
in each of these
areas!!

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



18

Stakeholder Tasks

Department	Tasks
Urogynecology Clinic: Management Schedulers and RNs	-Changes to clinic template -Updates with surgical pre/postop process workflow -Nutrition survey
Preop RN Staff	-Medication administration
PACU RN Staff	-Early mobilization, PO intake
Inpatient RN Staff	-Early mobilization, PO intake -Multimodal pain regimen
Anesthesia	-Identify multimodal pain regimen -Assist with CHO loading timing, fluid management

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

19



Stakeholder Tasks

Department	Tasks
Pharmacy	-Identify current narcotic use in Gyn surgery -Engage in changes with pre/postop Rx's
Nutrition	-Provide recommendations for malnutrition survey -Assist with CHO loading
EMR	-Edit preoperative and postoperative order sets
Patient Education	-Patient education videos

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

20



Protocol Implementation

- **Preoperative / Clinical**
- Preoperative Education and psychological preparation:
 - Provide optional tours of the facility
 - Booklet provided created with input from ALL stakeholders
 - Initiate discharge education and expectations

Tell them they're going home!

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

21



Avera Urogynecology ERAS packet

Getting Your Skin Ready for Surgery with Hibiclens®, Chlorhexidine gluconate (CHG) Soap-Adult

Incentive Spirometry
Improves Lung Function

Deep Vein Thrombosis (DVT) and Pulmonary Embolism (PE)
Prevention and Treatment

Smoking and Wound Healing

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

22



UROGYNECOLOGY ERAS PATIENT CHECK LIST		
Check Date/Time	TOPIC	ACTION
	Before Surgery	
		- Meet with your primary care provider (PCP) within 30 days prior to surgery. This is done to make sure you have a safe surgery. - Have lab work done, which will be ordered by your PCP. This must be done 14 days prior to surgery.
	The Week Before Surgery	
		- We will call 3-5 days prior to surgery to go over pre-operative instructions (medications, restrictions, etc.) and review necessary surgery information. - Avoid shaving the surgical area. - Eat a healthy diet that is high in protein. We may recommend nutritional supplements (Ensure or Boost drink) starting a few days before surgery. - The night before surgery, shower with Hibiclens® wash and follow the instructions in your folder. - The night before surgery eat a normal meal up until midnight. We recommend you drink 20 ounces of Ensure Pre-Surgery drink (two 10 oz bottles or 32 ounces Gatorade or apple juice without pulp the night prior to surgery. Do NOT do this if you are a Type 1 diabetic or have gastroperesis.) (avoid "diet" juices). If you have Type 2 Diabetes, recommend you discuss preoperative carbohydrate drinks with your primary doctor or endocrinologist. - Try to empty your bowels before surgery. You can use your normal laxatives and bowel medications, but maintain adequate fluid intake. Avoid laxatives the day of surgery.

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

23



The Morning of Surgery	
	- Take medications as discussed at time of pre-op call. - The morning of surgery, use the Hibiclens® medicated wash, following instructions in your folder. - Drink 1 (10 oz) bottle of Ensure Pre-Surgery drink or 20-40 ounces of Gatorade or apple juice without pulp 3 hours prior to your surgery time. Try and drink this within 5-10 minutes. (Do NOT do this if you are a Type 1 diabetic or have gastroperesis). - It is okay to have clear liquids up until your arrival time. (NO milk products.) - If you have Type 1 Diabetes, we encourage increased water intake up to 3 hours prior to procedure time. - If you have Type 2 Diabetes, we recommend you discuss preoperative carbohydrate drinks with your primary doctor or endocrinologist. - If you have Type 1 or 2 Diabetes, avoid "diet" juices as these do not provide the right amounts of carbohydrates. - It is ok to have a light meal 4 hours prior to arrival time (non-fatty foods - dry toast & clear liquids). Do NOT eat 8 hours prior to scheduled arrival time (fried/fatty foods, meat, or large meals). - Arrival time will be given to you during the preoperative call.
In the Pre-Anesthesia Room	
	- You will change into a hospital gown. - An IV will be placed and labs may be drawn. - You will be given medications before the surgery to help with post-operative pain and nausea. You are allowed to take these with sips of water.
In the Operating Room	
Oxygen	If needed, oxygen may be used while in the operating room.
Pain Management	- Take prescribed medications to help with your post-op pain. If this does not alleviate your pain, tell a nurse. IV pain medications will also be used. - There may be some pressure in the vagina following surgery as a vaginal packing is often used to help decrease bleeding risks.
Nausea Management	- Sometimes a scopolamine patch is placed the night before or AM of surgery and will stay on for a total of 72 hours. - You will be given medications to help with nausea in the hospital.
Urinary Management	You may have a Foley catheter in place after your surgery.

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

24



In the Hospital	
Mobility	- You will be helped out of bed after surgery. - You will walk a minimum of four times per day. - You will be up in the chair for all meals.
Diet	- We will encourage a full diet and early initiation of drinking fluids. - Once you are adequately drinking fluids we will stop your IV. - Chew and swallow many baby bites your bowel function returns to normal quickly.
Pain Management	- Your pain should be well-controlled with oral medicines as soon as possible. - Ask your healthcare team questions about pain before you go home. - When you are in pain it is sometimes uncomfortable to take deep breaths. We encourage you to use Incentive Spirometry device to help your lung function. - Sometimes a scopolamine patch is placed the night before or AM of surgery and will stay on for a total of 72 hours.
Nausea Management	- You will be given medications to help with nausea in the hospital. - You may go home with a Foley catheter or have what is called a voiding trial. A voiding trial helps us know if you need to go home with a Foley catheter. - If vaginal packing was placed, it will be removed the following morning.
Urinating After Surgery	

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

Discharging From the Hospital	
Pain Management	You will be discharged home with pain medications.
Diet & Nausea	- Review the information on diet and nausea in your surgical folder. - We may prescribe medications for nausea if necessary for you to use at home.
Discharge Planning	You are ready to be discharged once you can walk, eat and drink, tolerate oral pain medication, and have the surgeon's approval. This may be the same day as your surgery OR the day after.
Discharge Instructions	- Review discharge instructions with your nurse before you go home. - Do not push, pull, or lift more than 10 lbs. for 6 weeks. - You can walk and go up the stairs, but do not overexert yourself. - Avoid intercourse, hot tubs, swimming pools, and hot baths for 6 weeks.
Catheter Management	- If you cannot urinate and have a Foley catheter at the time of discharge, the nurse will schedule an appointment for you to have a voiding trial. - The nurse will teach you how to care for your indwelling catheter, including how to plug it and connect the bag to it.
Constipation	- We will give you Miralax stool softeners to keep your bowels regular. - It is ok to resume your normal bowel medications, unless directed otherwise by your provider.
Activity	We encourage you to be active after surgery. Review the instructions in your folder and we will review any specific activities or answer any questions you have about specific activity before you are discharged.
Follow-up	- The nurses will give you a follow-up appointment in our clinic. Follow up in one week if you go home with a Foley catheter. Otherwise plan on following up in two weeks for a post-op check. - Dr. Barker will call you on the Friday following your surgery to check in.

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

Prehabilitation

- 1. Aerobic and resistance exercises
 - 2. Targeted functional exercise
 - 3. Dietary interventions
 - 4. Psychological interventions to reduce stress
- COMING SOON
- Patients with impaired pre-operative function likely gain greatest clinical benefit

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

Protocol Implementation

- Inpatient Procedures**
- Week prior: If positive nutrition screen → Nutrition consult
 - Use high-protein drinks the week before surgery
 - Phone call from Preop RNs
- Day prior:
 - Bowel prep only when necessary
 - Carbohydrate loading drink

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

Protocol Implementation

- Day of Surgery**
 - May ingest fluids until **4 hours** prior to anesthesia
 - Alternate literature: Light meal **6 hours** prior to anesthesia, with clear liquids up to **2 hours** prior to anesthesia



AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

Protocol Implementation

- Day of Surgery**
- Preoperative medications

Celecoxib	400 mg PO once
Gabapentin	600 mg PO once
Acetaminophen	1000 mg PO once

- Glucose POC <200
- Urogynecology patients:
 - Ua Screen (+/-)

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

Surgical Site Infection Reduction

- Antimicrobial prophylaxis
- Skin preparation
- Avoiding hypothermia
- Avoiding surgical drains
- Reducing perioperative hyperglycemia



AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

31

Avera
Medical Group
Urogynecology

Protocol Implementation

• Intraoperative Steps to Prevent PONV

- Nausea & Vomiting
 - Risk factors= female gender, gynecologic surgery, MIS
 - Transdermal Scopolamine patch
 - Decadron 4mg IV
 - Ondansetron 4-8mg IV

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

32

Avera
Medical Group
Urogynecology

Protocol Implementation

• Intraoperative

- Maintain Intraoperative Euvolemia:
- Decrease crystalloid administration
 - 0.9% Normal Saline, D5 ½ Normal Saline
- Increase colloid administration if needed
 - PRBCs, FFP, Albumin, Dextran, etc.

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

33

Avera
Medical Group
Urogynecology



AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

34

Avera
Medical Group
Urogynecology

Protocol Implementation

• Postoperative

- Limited foley placement
- Remove intraoperative if possible
 - Backfill bladder with 150cc prior to intraop removal
 - Discharged 1 hour sooner, \$400 cost savings/patient
- If foley is left in place, perform **retrograde active voiding trial**
 - Faster time to decision for discharge and higher void rate



AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

35

Avera
Medical Group
Urogynecology

Retrograde Voiding Trial

- Backfill foley with 300 cc sterile water and clamp
- Discontinue foley catheter
- Patient has 15 minutes to void >150 cc
- If unable, replace foley for discharge
- Return to office in 24-48 hours for removal
- NO BLADDER SCANNING



AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110

36

Avera
Medical Group
Urogynecology

Protocol Implementation

- **Postoperative**
- Multimodal pain regimen

Oxycodone	5-15 mg PO q 4 hours
Ibuprofen	600 mg PO q 6 hours
Acetaminophen	1000 mg PO q 8 hours
Docusate Sodium	100 mg PO q 12 hours
Miralax	17 g daily prn

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



37

Protocol Implementation

- **Postoperative**
- Accelerated ambulation
 - POD 0: out of bed >2 hours (>= 1 walk and sitting in chair)
 - POD 1: out of bed >8 hours (>= 4 walks and sitting in chair)
 - Up in chair for ALL meals

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



38

Protocol Implementation

- **Postoperative Diet**
- No OG Tube
 - If used in OR, remove at extubation
- IVF 40cc/hr until 600 cc fluid consumed
- Low residual diet starting 4h post surgery
- POD 0: Oral intake by midnight 800-2000 ml
- POD 1: 1500-2000 ml of fluids
- Osmotic diuretics: senna and docusate sodium, magnesium oxide prn

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



39

Protocol Implementation

- **Postoperative**
- Postoperative patient education
 - Provided by PACU/floor nurses
 - Consistent with information provided in clinic
 - Discharge instructions provided by providers
- I call all my postoperative patients within first week
- Rxs provided preoperatively

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



40

Predictive Calculator for Postop Opioid Use

1. Do you have a history of chronic pelvic pain?	No (0)	Yes (10)			
2. Do you have a history of endometriosis?	No (0)	Yes (10)			
3. Do you currently use any opioid pain medication regularly?	No (0)	Yes (10)			
4. On an average day, how much pain do you experience on a scale of 0 to 10? (0= no pain, 10= worst pain imaginable)	No pain (0)	Worst pain imaginable (10)			
5. How much pain do you expect to experience after your surgery on a scale from 0 to 10? (0= no pain, 10= worst pain imaginable)	No pain (0)	Worst pain imaginable (10)			
6. What do you expect your pain medication requirements will be after surgery? (0= no pain medication needed, 10= highest possible amount of pain medication needed)	No pain medication needed (0)	Highest possible amount of pain medication needed (10)			
7. Over the past 2 weeks, how often have you been bothered by little interest or pleasure in doing things?	Not at all (0)	Several days (2)	More than half the days (4)	Nearly every day (6)	
8. Over the past two weeks, how often have you been bothered by nothing but much about different things?	Not at all (0)	Several days (2)	More than half the days (4)	Nearly every day (6)	
9. To what degree do you have these thoughts and feelings when you are experiencing pain? When I am in pain, it's awful and I feel that I can't believe it.	Not at all (0)	To a slight degree (2)	To a moderate degree (4)	To a great degree (6)	All the time (8)
Total					

If score >= 33: Increased risk of using >15 tablets of oxycodone 5mg

If score <12: Likely to use <= 5 tablets of oxycodone 5mg

Wong M et al. AIOG 2019

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



41



- Open hysterectomy 0-20 tablets
- MIS hysterectomy 0-10 tablets
- Cesarean section 0-10 tablets
- Vaginal delivery 0 tablets

Overton et al. Am Coll Surg 2018

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



42

Patient Education Video

- <https://avera-51.wistia.com/medias/39a5bkn18u>

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



43

Successes and Struggles

Successes

- Shorter length of stay
- High patient satisfaction scores
- No change in complications or readmissions

Struggles

- Difficulties getting EMR to add nutrition screen, order sets
- Coordinating all key stakeholders to incorporate changes
- Gauging adequate postoperative narcotic tablets

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



44

Key Points

- Implementing ERAS can have a positive impact on shorter length of stay, quicker ambulation, quicker return to bowel function, cost savings, and patient satisfaction
- Incorporating key stake holders in ERAS implementation is pivotal in successful role out and implementation

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



45

Questions?

- Does ERAS decrease the risk of complications in surgery? True or False
- Glycemic control is **NOT** important for minimizing complications during the perioperative period for surgical patients? True or False

AveraUrogyn.org | 605-504-1900
6100 S. Louise Ave, Suite 1110



46