



YOU ARE INVITED TO AN ALEXION-SPONSORED Neuromyelitis Optica Spectrum Disorder (NMOSD) Speaker Program

INDICATION

ULTOMIRIS is indicated for the treatment of adult patients with neuromyelitis optica spectrum disorder (NMOSD) who are anti-aquaporin-4 (AQP4) antibody positive.

Ultomiris NMOSD Abbreviated Branded Speaker Deck

Warren Felton III, MD
VCU Health System

Thursday, April 2, 2026

6:00 PM EST

Maggiano's Little Italy

11800 West Broad Street
Richmond, VA 23233

Limited seating available

RSVP by 3/27/2026

Registrations may still be accepted after this date pending availability

RSVP

Please contact
Edmund Moore
at (973) 865-7660
or edmund.moore@alexion.com

Please refer to meeting ID number: 18073

Select Important Safety Information, Including Boxed WARNING

WARNING: SERIOUS MENINGOCOCCAL INFECTIONS

ULTOMIRIS, a complement inhibitor, increases the risk of serious infections caused by *Neisseria meningitidis* [see *Warnings and Precautions (5.1)*] Life-threatening and fatal meningococcal infections have occurred in patients treated with complement inhibitors. These infections may become rapidly life-threatening or fatal if not recognized and treated early.

Alexion is committed to complying with applicable laws and regulations and adhering to the highest standards in its interactions with healthcare professionals, including those that govern the provision of modest meals to attendees at company-sponsored events. Per industry guidelines, we are unable to accommodate spouses or guests at this event. The invitation is nontransferable and is only for individuals who have a professional interest in the educational information that will be shared at the event. In order to ensure accurate transparency reporting of meals, Alexion requires program attendees to sign in upon arrival. Subject to federal and state regulations, Alexion will disclose information related to meals provided to attendees, as applicable. In most cases, this information will be made public. Attendees may opt out of the meal by indicating so on their RSVP. Certain states and federal agencies have defined the maximum fair market value allowable for modest meals given to healthcare professionals or prohibited the receipt of meals at company-sponsored events. You are accountable for understanding such restrictions and complying with them. If you are impacted by those restrictions, we respectfully request that you not partake in the meal offered by Alexion at the event. Please ask the Alexion representative at the event for information about purchasing food at the event.

Please see additional Important Safety Information on pages 2 and 3 and the accompanying full [Prescribing Information](#) for ULTOMIRIS, including Boxed WARNING regarding serious and life-threatening or fatal meningococcal infections.



IMPORTANT SAFETY INFORMATION

ULTOMIRIS, a complement inhibitor, increases the risk of serious infections caused by *Neisseria meningitidis* [see *Warnings and Precautions (5.1)*] Life-threatening and fatal meningococcal infections have occurred in patients treated with complement inhibitors. These infections may become rapidly life-threatening or fatal if not recognized and treated early.

- Complete or update vaccination for meningococcal bacteria (for serogroups A, C, W, Y, and B) at least 2 weeks prior to the first dose of ULTOMIRIS, unless the risks of delaying ULTOMIRIS therapy outweigh the risk of developing a serious infection. Comply with the most current Advisory Committee on Immunization Practices (ACIP) recommendations for vaccinations against meningococcal bacteria in patients receiving a complement inhibitor. See *Warnings and Precautions (5.1)* for additional guidance on the management of the risk of serious infections caused by meningococcal bacteria.
- Patients receiving ULTOMIRIS are at increased risk for invasive disease caused by *Neisseria meningitidis*, even if they develop antibodies following vaccination. Monitor patients for early signs and symptoms of serious meningococcal infections and evaluate immediately if infection is suspected.

Because of the risk of serious meningococcal infections, ULTOMIRIS is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS) called ULTOMIRIS and SOLIRIS REMS [see *Warnings and Precautions (5.2)*].

CONTRAINDICATIONS

- Initiation in patients with unresolved serious *Neisseria meningitidis* infection.

WARNINGS AND PRECAUTIONS

Serious Meningococcal Infections

ULTOMIRIS, a complement inhibitor, increases a patient's susceptibility to serious, life-threatening, or fatal infections caused by meningococcal bacteria (septicemia and/or meningitis) in any serogroup, including non-groupable strains. Life-threatening and fatal meningococcal infections have occurred in both vaccinated and unvaccinated patients treated with complement inhibitors.

Revaccinate patients in accordance with ACIP recommendations considering the duration of ULTOMIRIS therapy. Note that ACIP recommends an administration schedule in patients receiving complement inhibitors that differs from the administration schedule in the vaccine prescribing information. If urgent ULTOMIRIS therapy is indicated in a patient who is not up to date with meningococcal vaccines according to ACIP recommendations, provide antibacterial drug prophylaxis and administer meningococcal vaccines as soon as possible. Various durations and regimens of antibacterial drug prophylaxis have been considered, but the optimal durations and drug regimens for prophylaxis and their efficacy have not been studied in unvaccinated or vaccinated patients receiving complement inhibitors, including ULTOMIRIS. The benefits and risks of treatment with ULTOMIRIS, as well as those associated with antibacterial drug prophylaxis in unvaccinated or vaccinated patients, must be considered against the known risks for serious infections caused by *Neisseria meningitidis*.

Vaccination does not eliminate the risk of serious meningococcal infections, despite development of antibodies following vaccination.

Closely monitor patients for early signs and symptoms of meningococcal infection and evaluate patients immediately if infection is suspected. Inform patients of these signs and symptoms and instruct patients to seek immediate medical care if they occur. Promptly treat known infections. Meningococcal infection may become rapidly life-threatening or fatal if not recognized and treated early. Consider interruption of ULTOMIRIS in patients who are undergoing treatment for serious meningococcal infection depending on the risks of interrupting treatment in the disease being treated.

Please see additional Important Safety Information on next page and the accompanying full [Prescribing Information](#) for ULTOMIRIS, including Boxed WARNING regarding serious and life-threatening or fatal meningococcal infections.

ALEXION, the Alexion logo, ULTOMIRIS, and SOLIRIS are registered trademarks of Alexion Pharmaceuticals, Inc.
© 2024, Alexion Pharmaceuticals, Inc. All rights reserved. US/ULT-N/0145 V3 11/2024



ULTOMIRIS and SOLIRIS REMS

Due to the risk of serious meningococcal infections, ULTOMIRIS is available only through a restricted program called ULTOMIRIS and SOLIRIS REMS.

Prescribers must enroll in the REMS, counsel patients about the risk of serious meningococcal infection, provide patients with the REMS educational materials, assess patient vaccination status for meningococcal vaccines (against serogroups A, C, W, Y, and B) and vaccinate if needed according to current ACIP recommendations two weeks prior to the first dose of ULTOMIRIS. Antibacterial drug prophylaxis must be prescribed if treatment must be started urgently, and the patient is not up to date with both meningococcal vaccines according to current ACIP recommendations at least two weeks prior to the first dose of ULTOMIRIS. Patients must receive counseling about the need to receive meningococcal vaccines and to take antibiotics as directed, signs and symptoms of meningococcal infection, and be instructed to carry the Patient Safety Card at all times during and for 8 months following ULTOMIRIS treatment.

Further information is available at www.UltSolREMS.com or [1-888-765-4747](tel:1-888-765-4747).

Other Infections

Serious infections with *Neisseria* species (other than *Neisseria meningitidis*), including disseminated gonococcal infections, have been reported.

ULTOMIRIS blocks terminal complement activation; therefore, patients may have increased susceptibility to infections, especially with encapsulated bacteria, such as infections caused by *Neisseria meningitidis* but also *Streptococcus pneumoniae*, *Haemophilus influenzae*, and to a lesser extent, *Neisseria gonorrhoeae*. Patients receiving ULTOMIRIS are at increased risk for infections due to these organisms, even if they develop antibodies following vaccination.

Thromboembolic Event Management

The effect of withdrawal of anticoagulant therapy during treatment with ULTOMIRIS has not been established. Treatment should not alter anticoagulant management.

Infusion-Related Reactions

Administration of ULTOMIRIS may result in systemic infusion-related reactions, including anaphylaxis and hypersensitivity reactions. In clinical trials, infusion-related reactions occurred in approximately 1 to 7% of patients, including lower back pain, abdominal pain, muscle spasms, drop or elevation in blood pressure, rigors, limb discomfort, drug hypersensitivity (allergic reaction), and dysgeusia (bad taste). These reactions did not require discontinuation of ULTOMIRIS. If signs of cardiovascular instability or respiratory compromise occur, interrupt ULTOMIRIS and institute appropriate supportive measures.

ADVERSE REACTIONS

Most common adverse reactions in adult patients with NMOSD (incidence $\geq 10\%$) were COVID-19, headache, back pain, arthralgia, and urinary tract infection. Serious adverse reactions were reported in 8 (13.8%) patients with NMOSD receiving ULTOMIRIS.

DRUG INTERACTIONS

Plasma Exchange, Plasmapheresis, and Intravenous Immunoglobulins

Concomitant use of ULTOMIRIS with plasma exchange (PE), plasmapheresis (PP), or intravenous immunoglobulin (IVIg) treatment can reduce serum ravulizumab concentrations and requires a supplemental dose of ULTOMIRIS.

Neonatal Fc Receptor Blockers

Concomitant use of ULTOMIRIS with neonatal Fc receptor (FcRn) blockers (e.g., efgartigimod) may lower systemic exposures and reduce effectiveness of ULTOMIRIS. Closely monitor for reduced effectiveness of ULTOMIRIS.

USE IN SPECIFIC POPULATIONS

Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to ULTOMIRIS during pregnancy. Healthcare providers and patients may call [1-833-793-0563](tel:1-833-793-0563) or go to www.UltomirisPregnancyStudy.com to enroll in or to obtain information about the registry.

To report SUSPECTED ADVERSE REACTIONS, contact Alexion Pharmaceuticals, Inc. at [1-844-259-6783](tel:1-844-259-6783) or FDA at [1-800-FDA-1088](tel:1-800-FDA-1088) or www.fda.gov/medwatch.

Please see full [Prescribing Information](#) for ULTOMIRIS, including Boxed WARNING regarding serious and life-threatening or fatal meningococcal infections.

ALEXION, the Alexion logo, ULTOMIRIS, and SOLIRIS are registered trademarks of Alexion Pharmaceuticals, Inc.

© 2024, Alexion Pharmaceuticals, Inc. All rights reserved. US/ULT-N/0145 V3 11/2024

