

Pain in Older Adults

International Association for the Study of Pain



We are facing a rapidly aging demographic across the world and pain prevalence is known to be the highest in this cohort, with one exception [2]. Recent systematic reviews of survey studies with quite large sample sizes support the notion that pain increases with advancing age. All studies support the concept that females are more prone to pain than males. The most common pain complaints were knees, hips, and low back pain. There was also a consensus that most pain was of musculoskeletal origin (osteoporosis or osteoarthritis) [13]. Aging and disability increases the potential for chronic pain [8]. The common pain sites are knees, hips, and low back often associated with osteoarthritis and osteoporosis. Females are more likely to develop chronic pain and it is often associated with obesity (McCarthy et al 2009, Patel et al 2013). Taken together, the increased risk for suffering from bothersome pain coupled with the reduced capacity to cope and avoid the potential harm(s) associated with pain highlights the special vulnerabilities of older segments of our community. A high incidence of neuropathic pain has been found in the nursing home population [15]. In aggregate, this situation represents a markedly increased risk for suffering from bothersome pain.

Pain and suffering often make the afflicted individual more vulnerable and this is especially true in the case of older adults. However, advanced age by itself can also lead to greater vulnerability potentially placing this segment of our population in double jeopardy. Older adults are known to have the highest incidence of disease; many of which can be painful [3]. Rates of surgery, procedural interventions, injury [1] and hospitalization are also highest in this age group [12]. Aging is often associated with slower healing and poorer recovery from acute injury or disease and this may result in a potentially greater risk of developing an ongoing, persistent pain problem [10].

Another important aspect of vulnerability relates to the potential for greater harm(s) in response to a precipitating event or condition. For a proportion of the older population, psychiatric (especially dementia) and medical comorbidity, frailty and loss of physiologic reserve may all decrease the capacity of the older individual to effectively deal with the negative aspects of untreated pain. Polypharmacy and comorbid disease may also reduce the number and type of available treatment options and so compromise effective management of bothersome pain [7]. For instance, 63% of older adults with dementia had bothersome chronic pain compared to 54% of adults without dementia in a sample of 7609 community dwelling older adults [5]. The relative lack of dedicated age-specific pain treatment programs, the lack of appropriate research on identifying age differences in pain and its impacts, as well as a long-recognized lack of randomized controlled trials conducted specifically in older populations has been noted [9]. As a result, there is a paucity of evidence to help guide current clinical practice and consequently a greater likelihood of harm in those older persons with problematic pain. A number of



© Copyright 2019 International Association for the Study of Pain. All rights reserved.

IASP brings together scientists, clinicians, health-care providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.

papers have discussed self-management of pain in this age cohort [6,14] primarily due to the lack of available pharmacological options.

In spite of our increasing awareness of the prevalence of pain amongst the older population and our understanding of the impact of pain upon this group, under-treatment remains prevalent. Widely held misconceptions held by health professionals and older adults themselves presents a barrier to adequate treatment. An interesting paper by Thielke et al (2012) identified four commonly held myths about pain and aging including: pain as a natural part of getting older; pain worsens over time; stoicism leads to pain tolerance; prescription analgesics are highly addictive. The paper reviewed the evidence behind each of these myths and concluded that pain is not a natural part of aging and actually remains stable over time. The fact that older adults are often stoical does not mean that they “get used” to pain. They also demonstrated that more than 80% of older adults with osteoarthritis wanted more information about the course of the disease, but about only one third had received this information.

So where do we go from here? We understand the issues around pain in older adults and we know that there is a high incidence of pain in this population, which is often confounded by communication barriers and misconceptions amongst health professionals. We need to find a way to educate our patients and our peers to understand these issues and look to manage pain more effectively for this population, perhaps working with them to help them to understand that they do not have to live with the pain, it is “not an expected part of aging”.

REFERENCES

- [1] Stubbs B, Eggermont L, Binnekade T, Sephery A, Patchay S, Schofield P. (2013) Pain and the risk for falls in community dwelling older adults: A systematic review and Meta-analysis . Archives of Physical Medicine and Rehabilitation [10 Sep 2013, 95(1):175-187. e9].
- [2] Fejer R, Ruhe A (2012) What is the prevalence of musculoskeletal problems in the elderly population in developed countries? A systematic critical literature review. *Chiropr Man Therap.* 2012; 20: 31.
- [3] Ferrucci, L Giallauria, F & Guralnik, J (2008) Epidemiology of Ageing. *Radiology Clinics of North America* July 46(4) 643- v
- [4] Hemmingsson ES, Gustafsson M, Isaksson U, Karlsson S, Gustafson Y, Sandman PO, Lövhelm H. (2018) Prevalence of pain and pharmacological pain treatment among old people in nursing homes in 2007 and 2013. *Eur J Clin Pharmacol.* 2018 Apr;74(4):483-488. doi: 10.1007/s00228-017-2384-2. Epub 2017 Dec 20.
- [5] Hunt LJ, Covinsky KE, Yaffe K, Stephens CE, Miao Y, Boscardin WJ, Smith AK. (2015) Pain in Community-Dwelling Older Adults with Dementia: Results from the National Health and Aging Trends Study. *J Am Geriatr Soc.* 2015 Aug;63(8):1503-11. doi: 10.1111/jgs.13536. Epub 2015 Jul 22.
- [6] Karttunen NM, Turunen JH, Ahonen RS, Hartikainen SA. (2015) Persistence of noncancer-related musculoskeletal chronic pain among community-dwelling older people: a population-based longitudinal study in Finland. *Clin J Pain.* 2015 Jan;31(1):79-85. doi: 10.1097/AJP.000000000000089.
- [7] Nobili A, Garattini S, Mannucci PM. Multiple diseases and polypharmacy in the elderly: challenges for the internist of the third millennium. *J Comorb.* 2011;1:28-44. Published 2011 Dec 27.
- [8] Molton I, Cook KF, Smith AE, Amtmann D, Chen WH, Jensen MP. Prevalence and impact of pain in adults aging with a physical disability: comparison to a US general population sample. *Clin J Pain.* 2014 Apr;30(4):307-15. doi: 10.1097/AJP.0b013e31829e9bca.
- [9] Reid MC, & Pillemer K. (2015) Management of chronic pain in older adults. *BMJ* 2015; 350
- [10] Schofield P (2007) Pain in Older Adults. *Rev Pain.* 2007 Aug; 1(1): 12–14
- [11] Smith AK, Cenzer IS, Knight SJ, Puntillo KA, Widera E, Williams BA, Boscardin WJ, Covinsky KE. (2010). The epidemiology of pain during the last 2 years of life. *Ann Intern Med.* 2010 Nov 2;153(9):563-9. doi: 10.7326/0003-4819-153-9-201011020-00005



© Copyright 2019 International Association for the Study of Pain. All rights reserved.

IASP brings together scientists, clinicians, health-care providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.

- [12] Søreide K, Wijnhoven. B (2016) Surgery for an Ageing Population. BJS 2016; 103: e7–e9
- [13] Woo J, Leung J, Lau E. (2009) Prevalence and correlates of musculoskeletal pain in Chinese elderly and the impact on 4-year physical function and quality of life. Public Health. 2009 Aug;123(8):549-56. doi: 10.1016/j.puhe.2009.07.006. Epub 2009 Aug 25
- Patel et al 2013
- [14] Tse M, Wan VT, Wong AM. (2013) Pain and pain-related situations surrounding community-dwelling older persons. J Clin Nurs. 2013 Jul;22(13-14):1870-9. doi: 10.1111/jocn.12238. Epub 2013 May 17
- [15] van Kollenburg EG, Lavrijsen JC, Verhagen SC, Zuidema SU, Schalkwijk A, Vissers KC. (2012) Prevalence, causes, and treatment of neuropathic pain in Dutch nursing home residents: a retrospective chart review. J Am Geriatr Soc. 2012 Aug;60(8):1418-25. doi: 10.1111/j.1532-5415.2012.04078.x. Epub 2012 Jul 12.

AUTHORS

Patricia Schofield, PhD, Co-Chair Global Year Task Force
Faculty of Health, Education, Medicine and Social Care
Abertay University
Dundee, United Kingdom

Stephen Gibson, PhD
National Ageing Research Institute
Melbourne, Australia

About the International Association for the Study of Pain®

IASP is the leading professional forum for science, practice, and education in the field of pain. [Membership is open to all professionals](#) involved in research, diagnosis, or treatment of pain. IASP has more than 7,000 members in 133 countries, 92 national chapters, and 24 Special Interest Groups.

As part of the Global Year Against Pain in the Most Vulnerable, IASP offers a series of Fact Sheets that cover specific topics related to pain in vulnerable populations. These documents have been translated into multiple languages and are available for free download. Visit www.iasp-pain.org/globalyear for more information.



© Copyright 2019 International Association for the Study of Pain. All rights reserved.

IASP brings together scientists, clinicians, health-care providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.