



ECONOMIC RESEARCH
FEDERAL RESERVE BANK OF ST. LOUIS
WORKING PAPER SERIES

Combining Forecasts From Nested Models

Authors	Todd E. Clark, and Michael W. McCracken
Working Paper Number	2008-037A
Creation Date	October 2008
Citable Link	https://doi.org/10.20955/wp.2008.037
Suggested Citation	Clark, T.E., McCracken, M.W., 2008; Combining Forecasts From Nested Models, Federal Reserve Bank of St. Louis Working Paper 2008-037. URL https://doi.org/10.20955/wp.2008.037

Published In	Oxford Bulletin of Economics and Statistics
Publisher Link	https://doi.org/10.1111/j.1468-0084.2009.00547.x

Federal Reserve Bank of St. Louis, Research Division, P.O. Box 442, St. Louis, MO 63166

The views expressed in this paper are those of the author(s) and do not necessarily reflect the views of the Federal Reserve System, the Board of Governors, or the regional Federal Reserve Banks. Federal Reserve Bank of St. Louis Working Papers are preliminary materials circulated to stimulate discussion and critical comment.

Combining Forecasts From Nested Models ^{*}

Todd E. Clark

Federal Reserve Bank of Kansas City

Michael W. McCracken

Board of Governors of the Federal Reserve System

September 2008

Abstract

Motivated by the common finding that linear autoregressive models often forecast better than models that incorporate additional information, this paper presents analytical, Monte Carlo, and empirical evidence on the effectiveness of combining forecasts from nested models. In our analytics, the unrestricted model is true, but a subset of the coefficients are treated as being local-to-zero. This approach captures the practical reality that the predictive content of variables of interest is often low. We derive MSE-minimizing weights for combining the restricted and unrestricted forecasts. Monte Carlo and empirical analyses verify the practical effectiveness of our combination approach.

JEL Nos.: C53, C52

Keywords: forecast combination, predictability, forecast evaluation

word count: 10,893 words

^{*} *Clark (corresponding author)*: Economic Research Dept.; Federal Reserve Bank of Kansas City; 925 Grand; Kansas City, MO 64198; todd.e.clark@kc.frb.org. *McCracken*: Board of Governors of the Federal Reserve System; 20th and Constitution N.W.; Mail Stop #61; Washington, D.C. 20551; michael.w.mccracken@frb.gov. We gratefully acknowledge excellent research assistance from Taisuke Nakata and helpful comments from anonymous referees, Jan Groen, David Hendry, Jim Stock, seminar participants at the Deutsch Bundesbank and Federal Reserve Bank of Kansas City, and participants at the Bank of England Workshop on Econometric Forecasting Models and Methods, the 2005 World Congress of the Econometric Society, NBER Summer Institute, and Stanford's SITE workshop on economic forecasting. The views expressed herein are solely those of the authors and do not necessarily reflect the views of the Federal Reserve Bank of Kansas City, Board of Governors, Federal Reserve System, or any of its staff.

1 Introduction

Forecasters are well aware of the so-called principle of parsimony: “simple, parsimonious models tend to be best for out-of-sample forecasting...” (Diebold (1998)). Although an emphasis on parsimony may be justified on various grounds, parameter estimation error is one key reason. In many practical situations, estimating additional parameters can raise the forecast error variance above what might be obtained with a simple model. Such is clearly true when the additional parameters have population values of zero. But the same can apply even when the population values of the additional parameters are non-zero, if the marginal explanatory power associated with the additional parameters is low enough. In such cases, in finite samples the additional parameter estimation noise may raise the forecast error variance more than including information from additional variables lowers it.¹

As this discussion suggests, parameter estimation noise creates a forecast accuracy tradeoff. Excluding variables that truly belong in the model could adversely affect forecast accuracy. Yet including the variables could raise the forecast error variance if the associated parameters are estimated sufficiently imprecisely. In light of such a tradeoff, combining forecasts from the unrestricted and restricted (or parsimonious) models could improve forecast accuracy. Such combination could be seen as a form of shrinkage, which various studies, such as Stock and Watson (2003), have found to be effective in forecasting.

For non-nested models, the motivation for model combination is clear even if the population values of the parameters are known; combination integrates the two distinct information sets being used in the models. Optimal weights are then a regression exercise (Bates and Granger, 1969). However, in the case of nested models, this approach does not work. If the population values of the parameters are known, one of the models necessarily forecast encompasses the other and hence the optimal combining weights are trivially either zero or one. Therefore, combination can only be relevant for nested models if the parameters are estimated and the sample size is finite. In such an environment, and under some simplifying assumptions such as strict exogeneity of regressors and i.i.d. errors, it is possible to work through one-step

¹For example, simulation evidence in Clark and McCracken (2006) shows that even though the true model relates inflation to the output gap, in finite samples a simple AR model for inflation will often (although not always) forecast as well as or better than the true model.

ahead forecast error variance calculations to determine the combining weights that would be optimal for forecasting in period $T + 1$, based on models estimated with T observations. However, such analytics are very limiting — ruling out, for example, lagged dependent variables and conditionally heteroskedastic errors.

Accordingly, this paper uses a different approach to develop a general theoretical basis for combining forecasts from nested models, and provides Monte Carlo and empirical evidence on the effectiveness of the proposed combinations. Our analytics are based on models we characterize as “weakly” nested: the unrestricted model is the true model, but a subset of the coefficients (those not part of the restricted model) are treated as being local-to-zero.² This analytic approach captures the practical reality that the predictive content of some variables of interest is often quite low. That the unrestricted model “converges” to the restricted model might, at face value, be seen as counterintuitive. However, the local asymptotics should be seen as a convenient analytical device, rather than a modeling procedure. This device allows us to capture the case in between the extremes noted above — that either the restricted model or the unrestricted model perfectly forecast encompasses the other. The same type of analytical device has been used effectively in the literatures on unit-root or near-unit root inference and weak instruments, despite limiting case implications that might also seem counterfactual (e.g., implying unit roots in inflation or interest rates, or instruments uncorrelated with endogenous variables).³

Under the weak nesting specification, we are able to derive weights for combining the forecasts from estimates of the restricted and unrestricted models that are optimal in the sense of minimizing the forecast mean square error (MSE). We then characterize the settings under which the combination forecast will be more accurate than the restricted or unrestricted forecasts. In the special case in which the coefficients on the extra variables in the unrestricted model are of a magnitude that makes the restricted and unrestricted models equally accurate, the MSE-minimizing forecast is a simple, equally-weighted average of the restricted and unrestricted forecasts.

In the Monte Carlo and empirical analysis, we show our proposed approach of

²Although we focus the presented analysis on nested linear models, our results could be generalized to nested nonlinear models.

³In fact, Hansen (2008) uses near-unit root asymptotics to motivate model averaging of OLS-estimated autoregressive models that either do (restricted) or do not (unrestricted) impose a unit root in much the same way we do using local-to-zero asymptotics.

combining forecasts from nested models to be effective for improving accuracy. Admittedly, the gains to averaging are often modest or even small. However, the gains are very consistent: in practice, in our results, averaging is very likely to improve on the accuracy of both the restricted and unrestricted model forecasts. Moreover, in practice, most of the benefits can be achieved at low cost, via simple, equal-weight averages. These simple averages typically perform at least as well as more complicated averages. In particular, in the applications, our proposed combination approaches work well compared to related alternatives, consisting of Bayesian-type estimation with priors that push certain coefficients toward zero and Bayesian model averaging of the restricted and unrestricted models.

Our results build on much prior work on forecast combination. Research focused on non-nested models ranges from the early work of Bates and Granger (1969) to recent contributions such as Stock and Watson (2003) and Elliott and Timmermann (2004). Combination of nested model forecasts has been considered only occasionally, in such studies as Goyal and Welch (2003) and Hendry and Clements (2004). Our approach most closely resembles the nested model combination in Hansen (2008) but for a stationary rather than non-stationary environment. Forecasts based on Bayesian model averaging as applied in such studies as Wright (2003) and Jacobson and Karlsson (2004) could also combine projections from nested models.

The paper proceeds as follows. Section 2 provides theoretical results on the possible gains from combination of forecasts from nested models. In section 3 we present Monte Carlo evidence on the finite sample effectiveness of our proposed forecast combination methods. Section 4 compares the effectiveness of the forecast methods in a range of empirical applications. Section 5 concludes. Additional theoretical details are presented in Clark and McCracken (2006).

2 Theory

We begin by using a simple example to illustrate our essential ideas and results. We then proceed to the more general case. After detailing the necessary notation and assumptions, we provide an analytical characterization of the bias-variance tradeoff, created by weak predictability, involved in choosing among restricted, unrestricted, and combined forecasts. We then derive the optimal combination weights.

2.1 A simple example

Suppose we are interested in forecasting y_{t+1} using a simple model relating y_{t+1} to a constant and a strictly exogenous, scalar variable x_t . Suppose, however, that the predictive content of x_t for y_{t+1} may be weak. To capture this possibility, we model the population relationship between y_{t+1} and x_t using local-to-zero asymptotics, such that, in large samples, the predictive content of x_t shrinks to zero (assume that, apart from the local element, the model fits in the framework of the usual classical normal regression model, with homoskedastic errors, etc.):

$$y_{t+1} = \beta_0 + T^{-1/2}\beta_1 x_t + u_{t+1}, \quad E(x_t u_{t+1}) = 0, \quad E(u_{t+1}^2) = \sigma^2. \quad (1)$$

In light of x 's weak predictive content, the forecast from an estimated model relating y_{t+1} to a constant and x_t (henceforth, the *unrestricted* model) could be less accurate than a forecast from a model relating y_{t+1} to just a constant (the *restricted* model). Whether that is so depends on the “signal” and “noise” associated with x_t and its estimated coefficient. Under the local asymptotics incorporated in the DGP (1), the signal-to-noise ratio is proportional to $\beta_1^2 \sigma_x^2 / \sigma^2$. Given σ^2 and σ_x^2 (or β_1), higher values of the coefficient on x (or the variance of x) raise the signal relative to the noise; given the other parameters, a higher residual variance σ^2 increases the noise, reducing the signal-to-noise ratio. In general, noise associated with estimating the coefficient on x creates a forecast accuracy tradeoff. Excluding x could adversely affect forecast accuracy, while including it could increase the forecast error variance if the coefficient is estimated sufficiently imprecisely.

In light of this tradeoff between predictive content and additional noise from parameter estimation, a combination of the unrestricted and restricted model forecasts could be more accurate than either of the individual forecasts. We consider a combined forecast that puts a weight of α_t^* on the restricted model forecast and $1 - \alpha_t^*$ on the unrestricted model forecast. We then analytically determine the weight α_t^* that yields the forecast with lowest expected squared error in period $t + 1$.

As we establish more formally below, the (estimated) MSE-minimizing combination weight α_t^* is a function of the signal-to-noise ratio:

$$\hat{\alpha}_t^* = \left[1 + \left(\sqrt{t} \hat{b}_1 \right)^2 \hat{\sigma}_x^2 / \hat{\sigma}^2 \right]^{-1}, \quad (2)$$

where \hat{b}_1 denotes the coefficient on x ($\sqrt{t}\hat{b}_1$ corresponds to an estimate of the local

population coefficient β_1), $\hat{\sigma}_x^2$ denotes the variance of x , and $\hat{\sigma}^2$ denotes the residual variance, all estimated at time t (for forecasting at $t + 1$).⁴ As this result indicates, if the predictive content of x is such that the signal-to-noise ratio equals 1, then $\hat{\alpha}_t^* = .5$: the MSE-minimizing forecast is a simple average.

Admittedly, the local-to-zero asymptotic implication that the true model converges to the restricted model might strike some as counterintuitive. However, we view the local-to-zero setup as a convenient analytical device, as opposed to a modeling device, which ultimately leads to model combination that matches up with intuition. This device allows us to capture the case in between the extremes provided by conventional asymptotics — those extremes being that either the restricted model or the unrestricted model forecast encompasses the other. Under the local approximation, for a given β_1 , the predictive content of x_t declines to zero as the sample size diverges. This approximation allows us to derive limiting forecast moments such that even though the larger model is the true one, it may or may not be more accurate than the smaller model — a result that conventional asymptotics applied to estimated models cannot deliver under general conditions. But this approximation shouldn't be taken to mean we intend to model the predictive content of x_t as declining as forecasting moves forward for a given data sample. Rather, in a practical setting, we view the value of β_1/\sqrt{T} as being fixed, which implies that, as the sample expands, the implicit β_1 is increasing. In turn, as the sample expands as forecasting moves forward in time, the predictive content of x_t rises, such that the optimal combination forecast gradually puts increasing weight on the unrestricted model — as intuition suggests should occur, and indeed does in our Monte Carlo and empirical results.

2.2 The general case: environment

In the general case, the possibility of weak predictors is modeled using a sequence of linear DGPs of the form (**Assumption 1**)

$$y_{T,j+\tau} = x'_{T,2,j}\beta_T^* + u_{T,j+\tau} = x'_{T,1,j}\beta_1^* + x'_{T,22,j}(T^{-1/2}\beta_{22}^*) + u_{T,j+\tau}, \quad (3)$$

$$Ex_{T,2,j}u_{T,j+\tau} \equiv Eh_{T,j+\tau} = 0 \text{ for all } j = 1, \dots, t, \quad t = T - P + 1, \dots, T,$$

⁴Clements and Hendry (1998) derive a similar result, for the combination of a forecast based on the unconditional mean and a forecast based on an AR(1) model without intercept, the model assumed to generate the data.

where P denotes the number of predictions considered. Note that we allow the dependent variable $y_{T,j+\tau}$, the predictors $x_{T,2,j}$ and the error term $u_{T,j+\tau}$ to depend upon T , the final forecast origin. We make this explicit in the notation to emphasize that as the overall sample size is allowed to increase in our asymptotics, this parameterization affects their marginal distributions. While this is obvious for $y_{T,j+\tau}$ it is also true for $x_{T,2,j}$ if lagged values of the dependent variable are used as predictors. As such, our analytical results are based upon assumptions made on the triangular array $\{\{y_{T,j}, x'_{T,2,j}\}_{j=1}^{T+\tau}\}_{T \geq 1}$.

For a fixed value of T , our forecasting agent observes the sequence $\{y_{T,j}, x'_{T,2,j}\}_{j=1}^t$ sequentially at each forecast origin $t = T - P + 1, \dots, T$. Forecasts of the scalar $y_{T,t+\tau}$, $\tau \geq 1$, are generated using a $(k \times 1, k = k_1 + k_2)$ vector of covariates $x_{T,2,t} = (x'_{T,1,t}, x'_{T,22,t})'$, linear parametric models $x'_{T,i,t}\beta_i^*$, $i = 1, 2$, and a combination of the two models, $\alpha_t x'_{T,1,t}\beta_1^* + (1 - \alpha_t)x'_{T,2,t}\beta_2^*$. The parameters are estimated using OLS (**Assumption 2**) and hence $\hat{\beta}_{i,t} = \arg \min_{\beta_i} t^{-1} \sum_{j=1}^{t-\tau} (y_{T,j+\tau} - x'_{T,i,j}\beta_i)^2$, $i = 1, 2$, for the restricted and unrestricted models, respectively.⁵ We denote the loss associated with the τ -step ahead forecast errors as $\hat{u}_{T,i,t+\tau}^2 = (y_{T,t+\tau} - x'_{T,i,t}\hat{\beta}_{i,t})^2$, $i = 1, 2$, and $\hat{u}_{T,W,t+\tau}^2 = (y_{T,t+\tau} - \alpha_t x'_{T,1,t}\hat{\beta}_{1,t} - (1 - \alpha_t)x'_{T,2,t}\hat{\beta}_{2,t})^2$ for the restricted, unrestricted, and combined, respectively.

The following additional notation will be used. Let $H_{T,i}(t) = (t^{-1} \sum_{j=1}^{t-\tau} x_{T,i,j} u_{T,j+\tau}) = (t^{-1} \sum_{j=1}^{t-\tau} h_{T,i,j+\tau})$, $B_{T,i}(t) = (t^{-1} \sum_{j=1}^{t-\tau} x_{T,i,j} x'_{T,i,j})^{-1}$, and $B_i = \lim_{T \rightarrow \infty} (E x_{T,i,j} x'_{T,i,j})^{-1}$ for $i = 1, 2$. For $U_{T,j} = (h'_{T,2,j+\tau}, \text{vec}(x_{T,2,j} x'_{T,2,j}))'$, let $V = \sum_{l=-\tau+1}^{\tau-1} \Omega_{11,l}$, where $\Omega_{11,l}$ is the upper block-diagonal element of Ω_l defined below. For any $(m \times n)$ matrix A with elements $a_{i,j}$ and column vectors a_j , let: $\text{vec}(A)$ denote the $(mn \times 1)$ vector $[a'_1, a'_2, \dots, a'_n]'$; $|A|$ denote the max norm; and $\text{tr}(A)$ denote the trace. Let $\sup_t = \sup_{T-P+1 \leq t \leq T}$ and let \Rightarrow denote weak convergence. Finally, we define a variable selection matrix and a coefficient vector that appears directly in our key combination results: $J = (I_{k_1 \times k_1}, 0_{k_1 \times k_2})'$ and $\delta = (0_{1 \times k_1}, \beta_{22}^{*'})'$.

To derive our general results, we need two more assumptions (in addition to our assumptions (1 and 2) of a DGP with weak predictability and OLS-estimated linear

⁵In the interest of brevity, throughout the paper we focus on the recursive forecasting scheme, under which the estimation sample expands as forecasting moves forward in time. However, our results extend to the rolling scheme, under which the estimation sample is held at the same size and rolled forward as forecasting moves ahead in time. In a rolling scheme context, the t in equations (2) and (5) becomes the size of the rolling estimation sample and the summands begin with the first period in the rolling sample rather than period 1.

forecasting models).

Assumption 3: (a) $T^{-1} \sum_{j=1}^{[rT]} U_{T,j} U'_{T,j-l} \Rightarrow r\Omega_l$, $\Omega_l = \lim_{T \rightarrow \infty} T^{-1} \sum_{t=1}^T E(U_{T,j} U'_{T,j-l})$ for all $l \geq 0$, (b) $\Omega_{11,l} = 0$ all $l \geq \tau$, (c) $\sup_{T-P+1 \leq 1, s \leq T} E|U_{T,s}|^{2q} < \infty$ for some $q > 1$, (d) $U_{T,j} - EU_{T,j} = (h'_{T,2,j+\tau}, \text{vec}(x_{T,2,j} x'_{T,2,j} - Ex_{T,2,j} x'_{T,2,j}))'$ is a zero mean triangular array satisfying Theorem 3.2 of De Jong and Davidson (2000).

Assumption 4: For $s \in (1 - \lambda_P, 1]$, (a) $\alpha_t \Rightarrow \alpha(s) \in [0, 1]$, (b) $\lim_{T \rightarrow \infty} P/T = \lambda_P \in (0, 1)$.

Assumption 3 imposes three types of conditions. First, in (a) and (c) we require that the observables, while not necessarily covariance stationary, are asymptotically mean square stationary with finite second moments. We do so in order to allow the observables to have marginal distributions that vary as the weak predictive ability strengthens along with the sample size but are ‘well-behaved’ enough that, for example, sample averages converge in probability to the appropriate population means. Second, in (b) we impose the restriction that the τ -step ahead forecast errors are MA($\tau - 1$). We do so in order to emphasize the role that weak predictors have on forecasting without also introducing other forms of model misspecification. Finally, in (d) we impose the high level assumption that, in particular, $h_{T,2,j+\tau}$ satisfies Theorem 3.2 of De Jong and Davidson (2000). By doing so we not only insure (results needed in Clark and McCracken (2006)) that certain weighted partial sums converge weakly to standard Brownian motion, but also allow ourselves to take advantage of various results pertaining to convergence in distribution to stochastic integrals.

Our final assumption is unique: we permit the combining weights to change with time. In this way, we allow the forecasting agent to balance the bias-variance tradeoff differently across time as the increasing sample size provides stronger evidence of predictive ability. Finally, we impose the requirement that $\lim_{T \rightarrow \infty} P/T = \lambda_P \in (0, 1)$ and hence the duration of forecasting is finite but non-trivial.

2.3 Theoretical results on the tradeoff

Our characterization of the bias-variance tradeoff associated with weak predictability is based on $\sum_{t=T-P+1}^T (\hat{u}_{T,2,t+\tau}^2 - \hat{u}_{T,W,t+\tau}^2)$, the difference in the (normalized) MSEs of the unrestricted and combined forecasts. Clark and McCracken (2006) provide a

general characterization of the tradeoff, in Theorem 1. But in the absence of a closed form solution for the limiting distribution of the loss differential (the distribution provided in Clark and McCracken (2006)), we proceed in this section to focus on the mean of this loss differential.

From the general case proved in Clark and McCracken (2006), we first establish the expected value of the loss differential, in the following corollary.

Corollary 1: $E \sum_{t=T-P+1}^T (\hat{u}_{T,2,t+\tau}^2 - \hat{u}_{T,W,t+\tau}^2) \rightarrow \int_{1-\lambda_P}^1 E\xi_W(s) = \int_{1-\lambda_P}^1 (1 - (1 - \alpha(s))^2) s^{-1} \text{tr}((-JB_1J' + B_2)V) ds - \int_{1-\lambda_P}^1 \alpha^2(s) \delta' B_2^{-1} (-JB_1J' + B_2) B_2^{-1} \delta ds.$

This decomposition implies that the bias-variance tradeoff depends on: (1) the duration of forecasting (λ_P), (2) the dimension of the parameter vectors (through the dimension of δ), (3) the magnitude of the predictive ability (as measured by quadratics of δ), (4) the forecast horizon (via V , the long-run variance of $h_{T,2,t+\tau}$), and (5) the second moments of the predictors ($B_i = \lim_{T \rightarrow \infty} (Ex_{T,i,t}x'_{T,i,t})^{-1}$).

The first term on the right-hand side of the decomposition can be interpreted as the pure “variance” contribution to the mean difference in the unrestricted and combined MSEs. The second term can be interpreted as the pure “bias” contribution. Clearly, when $\delta = 0$ and thus there is no predictive ability associated with the predictors $x_{T,22,t}$, the expected difference in MSE is positive so long as $\alpha(s) \neq 0$. Since the goal is to choose $\alpha(s)$ so that $\int_{1-\lambda_P}^1 E\xi_W(s)$ is maximized, we immediately reach the intuitive conclusion that we should always forecast using the restricted model and hence set $\alpha(s) = 1$. When $\delta \neq 0$, and hence there is predictive ability associated with the predictors $x_{T,22,t}$, forecast accuracy is maximized by combining the restricted and unrestricted model forecasts. The following corollary provides the optimal combination weight. Note that, to simplify notation in the presented results, from this point forward we omit the subscript T from the predictors.

By maximizing the arguments of the integrals in Corollary 1 that contribute to the average expected mean square differential over the duration of forecasting, we obtain the optimal combination weight, given in Corollary 2.

Corollary 2: The pointwise optimal combining weights satisfy

$$\alpha^*(s) = \left[1 + s \left(\frac{\beta'_{22} (Ex_{22,t}x'_{22,t} - Ex_{22,t}x'_{1,t} (Ex_{1,t}x'_{1,t})^{-1} Ex_{1,t}x'_{22,t}) \beta_{22}}{\text{tr}((-JB_1J' + B_2)V)} \right) \right]^{-1}. \quad (4)$$

This combining weight is decreasing in the marginal ‘signal to noise’ ratio

$$s\beta'_{22}(Ex_{22,t}x'_{22,t} - Ex_{22,t}x'_{1,t}(Ex_{1,t}x'_{1,t})^{-1}Ex_{1,t}x'_{22,t})\beta_{22}/tr((-JB_1J' + B_2)V).$$

As the marginal ‘signal’, $s\beta'_{22}(Ex_{22,t}x'_{22,t} - Ex_{22,t}x'_{1,t}(Ex_{1,t}x'_{1,t})^{-1}Ex_{1,t}x'_{22,t})\beta_{22}$, increases, we place more weight on the unrestricted model and less on the restricted one. Conversely, as the marginal ‘noise’, $tr((-JB_1J' + B_2)V)$, increases, we place more weight on the restricted model and less on the unrestricted model. In the special case in which the signal-to-noise ratio equals 1, the optimal combination weight is 1/2. Finally, as forecasting moves forward in time and the estimation sample (represented by s) increases, we place increasing weight on the unrestricted model.

A bit more algebra establishes the determinants of the size of the benefits to combination. The expected loss difference $E\xi_W^*(s)$ takes the easily interpretable form

$$\frac{tr((-JB_1J' + B_2)V)^2}{s(s\beta'_{22}(Ex_{22,t}x'_{22,t} - Ex_{22,t}x'_{1,t}(Ex_{1,t}x'_{1,t})^{-1}Ex_{1,t}x'_{22,t})\beta_{22} + tr((-JB_1J' + B_2)V))}.$$

With a 1-step horizon and conditionally homoskedastic errors, the numerator simplifies to $tr((-JB_1J' + B_2)V) = \sigma^2 k_2$. In either case, we expect the optimal combination to provide the most benefit when the marginal ‘noise’, $tr((-JB_1J' + B_2)V)$, is large or when the marginal ‘signal’, $s\beta'_{22}(Ex_{22,t}x'_{22,t} - Ex_{22,t}x'_{1,t}(Ex_{1,t}x'_{1,t})^{-1}Ex_{1,t}x'_{22,t})\beta_{22}$, is small. And again, we obtain the result that, as the estimation sample grows, any benefits from combination vanish as the parameter estimates become more accurate.

Note, however, that the term $\beta'_{22}(Ex_{22,t}x'_{22,t} - Ex_{22,t}x'_{1,t}(Ex_{1,t}x'_{1,t})^{-1}Ex_{1,t}x'_{22,t})\beta_{22}$ is a function of the local-to-zero parameters β_{22} . Moreover, note that these optimal combining weights are not presented relative to an environment in which agents are forecasting in ‘real time’. Therefore, for practical use, we suggest a transformed formula. Let \hat{B}_i and \hat{V} denote estimates of B_i and V , respectively, based on data through period t .⁶ If we let $T^{1/2}\hat{\beta}_{22}$ denote an estimate of the local-to-zero parameter β_{22}^* and set $s = t/T$, we obtain the following real time estimate of the pointwise optimal combining weight:

⁶We estimate B_i with $\hat{B}_i = (t^{-1} \sum_{j=1}^{t-\tau} x_{i,j}x'_{i,j})^{-1}$, where $x_{i,t}$ is the vector of regressors in the forecasting model (supposing the MSE stationarity assumed in the theoretical analysis). At a forecast horizon (τ) of one period, we estimate V using $\hat{V} = t^{-1} \sum_{j=1}^{t-\tau} \hat{u}_{1,j}^2 x_{2,j}x'_{2,j}$. At longer forecast horizons, we similarly compute V with the Newey–West estimator (again, using the residual from the restricted model) and $2(\tau-1)$ lags. In all cases, we use the restricted model residual in computing V , in light of the evidence in such studies as Godfrey and Orme (2004) that imposing such restrictions improves the small sample properties of heteroskedasticity-robust variances.

$$\hat{\alpha}_t^* = \left[1 + t \left(\frac{\hat{\beta}'_{22} \left(t^{-1} \sum_{j=1}^{t-\tau} x_{22,j} x'_{22,j} - (t^{-1} \sum_{j=1}^{t-\tau} x_{22,j} x'_{1,j}) \hat{B}_1 (t^{-1} \sum_{j=1}^{t-\tau} x_{1,j} x'_{22,j}) \right) \hat{\beta}_{22}}{tr((-J\hat{B}_1 J' + \hat{B}_2)\hat{V})} \right) \right]^{-1}. \quad (5)$$

The parameter estimates provide asymptotically mean unbiased estimates of the local-to-zero parameters on which our theoretical derivations (Corollary 2) are based. Nonetheless, our estimates of the local-to-zero parameters are not consistent. The local-to-zero asymptotics allow us to derive closed-form solutions for the optimal combination weights, but require knowledge of local-to-zero parameters for which we can obtain mean unbiased, but not consistent, estimates via OLS (and rescaling). We therefore simply use rescaled OLS magnitudes as estimates of the assumed local-to-zero values and subsequent optimal combining weights. Below we use Monte Carlo experiments and empirical examples to determine whether the estimated quantities perform well enough to be a valuable tool for forecasting.

Conceptually, our proposed combination (5) might be seen as a variant of a Stein rule estimator.⁷ With conditionally homoskedastic, 1-step ahead forecast errors, the signal-to-noise ratio in our combination coefficient $\hat{\alpha}_t$ is the conventional F -statistic for testing the null of coefficients of 0 on the x_{22} variables. With additional (and strong) assumptions of normality and strict exogeneity of the regressors, the F -statistic has a non-central F distribution, with a mean that is a linear function of the population signal-to-noise ratio. Based on that mean, the population-level signal-to-noise ratio can be alternatively estimated as F -statistic $- 1$. A combination forecast based on this estimate is exactly the same as the forecast that would be obtained by applying conventional Stein rule estimation to the unrestricted model.

This Stein rule result suggests an alternative estimate of the optimal combination coefficient α_t^* with potentially better small sample properties. Specifically, based on (i) the equivalence of the directly estimated signal-to-noise ratio and the conventional F -statistic result and (ii) the centering of the F distribution at a linear transform of the population signal-to-noise ratio, we might consider replacing the signal-to-noise

⁷Our optimal, but infeasible, combining weights are closely related to the minimum-MSE estimator provided in Theil (1971). Our results primarily differ in that we permit serially correlated and conditionally heteroskedastic errors, and don't require strict exogeneity of the regressors.

ratio estimate in (5) with the signal-to-noise ratio estimate less 1. However, under this estimation approach, the combination forecast could put a weight of more than 1 on the restricted model and a negative weight on the unrestricted. As a result, we might consider a truncation that bounds the weight between 0 and 1:

$$\hat{\alpha}_t^* = [1 + \max(0, \text{signal/noise} - 1)]^{-1}, \quad (6)$$

where the signal/noise term is the same as that in the baseline estimator (5). In light of potential concerns about the small sample properties of the estimator (5), we include a forecast combination based on (6) in our Monte Carlo and empirical analyses.

More generally, in cases in which the marginal predictive content of the x_{22} variables is small or modest, a simple average forecast might be more accurate than our proposed estimated combinations based on (5) or (6). With β_{22} coefficients sized such that the restricted and unrestricted models are nearly equally accurate, the population-level optimal combination weight will be close to 1/2. As a result, forecast accuracy could be enhanced by imposing a combination weight of 1/2 instead of estimating it, in light of the potential for noise in the combination coefficient estimate. A parallel result is well-known in the non-nested combination literature: simple averages are often more accurate than estimated optimal combinations (see, e.g., Smith and Wallis (2008)).

Of course, in our context, the optimal weight changes over time, rising as more data become available for model estimation, such that an optimal weight that starts out (or ends up) close to 1/2 might not end up (or start out) close to 1/2. In practice, however, our estimated weights change only very gradually over time. For example, in the DGP 3 experiment in Table 1 presented below, the theoretically optimal combination weight (for a forecast horizon of 1 period) declines from only 0.5 to 0.4 over the first 40 observations of the forecast sample. As a result, as long as the optimal combination weight starts out in the neighborhood of 1/2, a simple average is likely to do well in samples of common size, even though the optimal weight is gradually declining over the course of the sample.

Our proposed combination (5) might also be expected to have some relationship to Bayesian methods. In the very simple case of the example of section 2.1, the proposed combination forecast corresponds to a forecast from an unrestricted model

with Bayesian posterior mean coefficients estimated with a prior mean of 0 and variance proportional to the signal–noise ratio. More generally, our proposed combination could correspond to the Bayesian model averaging considered in such studies as Wright (2003), Koop and Potter (2004), and Stock and Watson (2005).⁸ In the more general case, there may be some prior that makes a Bayesian average of the restricted and unrestricted forecasts similar to the combination forecast based on (5). But the rationale for Bayesian averaging is quite different from the combination rationale developed in this paper. Bayesian averaging is generally founded on model uncertainty. In contrast, our combination rationale is based on the bias–variance tradeoff associated with parameter estimation error, in an environment without model uncertainty.

3 Monte Carlo Evidence

We use Monte Carlo simulations of several multivariate data-generating processes to evaluate the finite-sample performance of the combination methods described above. In these experiments, the DGPs relate the predictand y to lagged y and lagged x , with the coefficients on lagged x set at various values. Forecasts of y are generated with the combination approaches considered above. Performance is evaluated using simple summary statistics of the distribution of each forecast’s MSE: the average MSE across Monte Carlo draws and the probability of equaling or beating the restricted model’s forecast MSE.

3.1 Experiment design

In light of the considerable practical interest in the out-of-sample predictability of inflation (see, for example, Stock and Watson (1999, 2003), Atkeson and Ohanian (2001), Orphanides and van Norden (2005), and Clark and McCracken (2006)), we present results for DGPs based on estimates of quarterly U.S. inflation models. In particular, we consider models based on the relationship of the change in core PCE inflation to (1) lags of the change in inflation and the output gap, (2) lags of the change in inflation, the output gap, and food and energy price inflation, and (3) lags of the change in inflation and five common business cycle factors, estimated as in Stock and

⁸Indeed, in the scalar environment of Stock and Watson (2005), setting their weighting function to $t\text{-stat}^2/(1 + t\text{-stat}^2)$ yields our combination forecast.

Watson (2005). We consider various combinations of forecasts from an unrestricted model that includes all variables in the DGP to forecasts from a restricted model that takes an AR form.

For each experiment, we conduct 10,000 simulations. With quarterly data in mind, we evaluate forecast accuracy over forecast periods of various lengths: $P = 1, 20, 40$, and 80. In our baseline results, the size of the sample used to generate the first (in time) forecast at horizon τ is $80 - \tau + 1$ (the estimation sample expands as forecasting moves forward in time). In light of the potential for forecast combination to yield larger gains with smaller model estimation samples, we also report selected results for experiments in which the size of the sample used to generate the first (in time) forecast at horizon τ is $40 - \tau + 1$.

The first DGP, based on the empirical relationship between the change in core inflation (Δy_t) and the output gap ($x_{1,t}$), takes the form

$$\begin{aligned}\Delta y_t &= -.40\Delta y_{t-1} - .18\Delta y_{t-2} - .09\Delta y_{t-3} - .04\Delta y_{t-4} + b_{11}x_{1,t-1} + u_t \\ x_{1,t} &= 1.15x_{1,t-1} - .05x_{1,t-2} - .20x_{1,t-3} + v_{1,t} \\ \text{var} \begin{pmatrix} u_t \\ v_{1,t} \end{pmatrix} &= \begin{pmatrix} .72 & \\ .02 & .57 \end{pmatrix}.\end{aligned}\tag{7}$$

We consider experiments with two different settings of b_{11} , the x_1 coefficient, which corresponds to our theoretical construct β_{22}/\sqrt{T} . The baseline value of b_{11} is the one that, in population, makes the null and alternative models equally accurate (in expectation, at the 1-step ahead horizon) in the first forecast period, period $T - P + 2$. Given the population moments implied by the DGP parameterization, this value is $b_{11} = .042$. The second setting we consider is the empirical value: $b_{11} = .10$.

The second DGP, based on estimated relationships among inflation (Δy_t), the output gap ($x_{1,t}$), and food and energy price inflation ($x_{2,t}$), takes the form:

$$\begin{aligned}\Delta y_t &= -.47\Delta y_{t-1} - .24\Delta y_{t-2} - .15\Delta y_{t-3} - .10\Delta y_{t-4} \\ &\quad + b_{11}x_{1,t-1} + b_{21}x_{2,t-1} + b_{22}x_{2,t-2} + u_t \\ x_{1,t} &= 1.15x_{1,t-1} - .05x_{1,t-2} - .20x_{1,t-3} + v_{1,t} \\ x_{2,t} &= .06x_{1,t-1} + .40x_{2,t-1} + .28x_{2,t-3} - .13x_{2,t-4} + v_{2,t} \\ \text{var} \begin{pmatrix} u_t \\ v_{1,t} \\ v_{2,t} \end{pmatrix} &= \begin{pmatrix} .62 & & \\ .03 & .57 & \\ -.06 & .06 & .70 \end{pmatrix}.\end{aligned}\tag{8}$$

As with DGP 1, we consider experiments with two settings of the set of b_{ij} coefficients, which correspond to the elements of β_{22}/\sqrt{T} . One setting is based on empirical estimates: $b_{11} = .07$, $b_{21} = .27$, $b_{22} = .10$. We take as the baseline experiment one in which all of these empirical values of the b_{ij} coefficients are multiplied by a constant less than one, such that, in population, the null and alternative models are expected to be equally accurate (at the 1-step ahead horizon) in (the first) forecast period $T - P + 2$. In our baseline experiments, this multiplying constant is .370.

The third DGP is based on estimated relationships among inflation (Δy_t) and five business cycle factors estimated as in Stock and Watson (2005) ($x_{i,t}, i = 1, \dots, 5$):

$$\begin{aligned}\Delta y_t &= -.40\Delta y_{t-1} - .19\Delta y_{t-2} - .10\Delta y_{t-3} - .04\Delta y_{t-4} + \sum_{i=1}^5 b_{i1}x_{i,t-1} + u_t \\ x_{i,t} &= \sum_{j=1}^4 a_{ij}x_{i,t-j} + v_{i,t}, \quad i = 1, \dots, 5.\end{aligned}\tag{9}$$

As with DGPs 1 and 2, we consider experiments with two different settings of the set of b_{ij} coefficients. One setting is based on empirical estimates: $b_{11} = .04$, $b_{21} = .09$, $b_{31} = .16$, $b_{41} = .04$, $b_{51} = .08$.⁹ We take as the baseline experiment one in which all of these empirical values of the b_{ij} coefficients are multiplied by a constant less than one, such that, in population, the null and alternative models are expected to be equally accurate (at the 1-step horizon) in forecast period $T - P + 2$. In our baseline experiments, this multiplying constant is .748.

3.2 Forecast approaches

Following practices common in the literature from which our applications are taken (see, e.g., Stock and Watson (2003)), direct multi-step forecasts one and four steps ahead are formed from various combinations of estimates of the following forecasting models:

$$y_{t+\tau}^{(\tau)} - y_t = \delta_0 + \sum_{j=1}^4 \delta_j \Delta y_{t+1-j} + u_{1,t+\tau}\tag{10}$$

$$y_{t+\tau}^{(\tau)} - y_t = \gamma_0 + \sum_{j=1}^4 \gamma_j \Delta y_{t+1-j} + \Gamma'_{22} x_{22,t} + u_{2,t+\tau},\tag{11}$$

⁹The coefficients of the AR models for the factors are as follows, in order from lags 1 to 4: factor 1: .81, -.18, .19, -.19; factor 2: .80, -.05, .16, -.18; factor 3: -.36, .16, .22, .12; factor 4: .31, .08, .39, .01; and factor 5: .25, .15, .24, .05. The residual variances of the five factors are as follows, in order for factors 1 through 5: 6.36, 2.35, .92, 2.08, 1.62. The variance of u_t is .67.

where $y_{t+\tau}^{(\tau)} = (1/\tau) \sum_{s=1}^{\tau} y_{t+s}$ and $y_{t+1}^{(1)} \equiv y_{t+1}$. In the actual inflation data underlying the DGP specification, $y_{t+\tau}^{(\tau)}$ corresponds to the average annual rate of price increase from period t to $t + \tau$. Across DGPs 1-3, the vector $x_{22,t}$ consists of, respectively, (1) $(x_{1,t})$, (2) $(x_{1,t}, x_{2,t}, x_{2,t-1})'$, and (3) $(x_{1,t}, x_{2,t}, x_{3,t}, x_{4,t}, x_{5,t})'$. Because the multi-step forecasts are projections of an average of y over the forecast horizon up to period $t + \tau$ rather than simply a projection of y in period $t + \tau$ (in order to follow the examples of the aforementioned studies), the relationship of forecast accuracy to horizon is unclear. Depending on the DGP, MSEs may rise or fall as the horizon increases.

We examine the accuracy of forecasts from: (i) OLS estimates of the restricted model (10); (ii) OLS estimates of the unrestricted model (11); (iii) the ‘known’ optimal linear combination of the restricted and unrestricted forecasts, using the weight implied by equation (4) and population moments implied by the DGP; (iv) the estimated optimal linear combination of the restricted and unrestricted forecasts, using the weight given in (5) and estimated moments of the data; (v) the estimated optimal linear combination using the Stein rule–variant weight given in (6); and (vi) a simple average of the restricted and unrestricted forecasts.¹⁰

3.3 Simulation results

In our Monte Carlo comparison of methods, we primarily base our evaluation on average MSEs over a range of forecast samples. For simplicity, in presenting average MSEs, we only report actual average MSEs for the restricted model (10). For all other forecasts, we report the ratio of a forecast’s average MSE to the restricted model’s average MSE. To capture potential differences in MSE distributions, we also present some evidence on the probabilities of equaling or beating the restricted model.

3.3.1 Results for signal = noise experiments

We begin with the case in which the coefficients b_{ij} (elements of β_{22}) on the lags of x_{it} (elements of x_{22}) in the DGPs (7)–(9) are set such that, at the 1-step ahead horizon, the restricted and unrestricted model forecasts for period $T - P + 2$ are expected to be equally accurate — because the signal and noise associated with the x_{it} variables are equalized as of that period. In this setting, the optimally combined forecast should, on

¹⁰As noted above, weights of 1/2 are optimal if the signal associated with the x variables equals the noise, making the models equally accurate.

average, be more accurate than either the restricted or unrestricted forecasts. Note, however, that the models are scaled to make only 1-step ahead forecasts equally accurate. At the 4-step ahead forecast horizon, the restricted model may be more or less accurate than the unrestricted, depending on the DGP.

The average MSE results reported in Table 1 confirm the theoretical implications. Consider first the 1-step ahead horizon. With all three DGPs, the ratio of the unrestricted model's average MSE to the restricted model's average MSE is close to 1.000 for all forecast samples. At the 4-step ahead horizon, for all DGPs the ratio of the unrestricted model's average MSE to the restricted model's average MSE is generally above 1.000. The unrestricted model fares especially poorly relative to the restricted in the case of DGP 3, in which the unrestricted model includes five more variables than the restricted. In general, in all cases, the MSE ratios for 4-step ahead forecasts from the unrestricted model tend to fall as P rises, reflecting the increase in the precision of the x coefficient (Γ_{22}) estimates that occurs as forecasting moves forward in time and the model estimation sample grows.

A combination of the restricted and unrestricted forecasts has a lower average MSE, with the gains generally increasing in the number of variables omitted from the restricted model and the forecast horizon. At the 1-step horizon, using the known optimal combination weight α_t^* yields $P = 20$ MSE ratios of .994, .983, and .974 for, respectively, DGPs 1, 2, and 3. At the 4-step horizon, the forecast based on the known optimal combination weight has $P = 20$ MSE ratios of .986, .962, and .973 for DGPs 1-3.

Not surprisingly, having to estimate the optimal combination weight tends to slightly reduce the gains to combination. For example, in the case of DGP 2 and $P = 20$, the MSE ratio for the estimated optimal combination forecast is .989, compared to .983 for the known optimal combination forecast. Using the Stein rule-based adjustment to the optimal combination estimate (based on equation (6)) has mixed consequences, sometimes faring a bit worse than the directly estimated optimal combination forecast (based on equation (5)) and sometimes a bit worse. To use the same DGP 2 example, the $P = 20$ MSE ratio for the Stein version of the estimated optimal combination is .990, compared to .989 for the directly estimated optimal combination. However, in the case of 4-step ahead forecasts for DGP 3 with the $P = 20$ sample, the MSE ratios of the known α_t^* , estimated $\hat{\alpha}_t^*$, and Stein-adjusted $\hat{\alpha}_t^*$ are, respectively,

.973, .991, and .985.

In the Table 1 experiments, the simple average of the restricted and unrestricted forecasts is consistently a bit more accurate than the estimated optimal combination forecast. For example, for DGP 3 and the $P = 20$ forecast sample, the MSE ratio of the simple average forecast is .974 for both 1-step and 4-step ahead forecasts, compared to the estimated optimal combination forecasts' MSE ratios of .982 (1-step) and .991 (4-step). There are two reasons a simple average fares so well. First, with the DGPs parameterized to make signal = noise for one-step ahead forecasts for period $T - P + 2$, the theoretically optimal combination weight is 1/2. Of course, as forecasting moves forward in time, the theoretically optimal combination weight declines, because as more and more data become available for estimation, the signal-to-noise ratio rises (e.g., in the case of DGP 3, the known optimal weight for the forecast of the 80th observation in the prediction sample is about .33). But the decline is gradual enough that only late in a long forecast sample would noticeable differences emerge between the theoretically optimal combination forecast and the simple average. Second, in practice, the optimal combination weight may not be estimated with much precision. As a result, imposing a fixed weight of 1/2 is likely better than trying to estimate a weight that is not dramatically different from 1/2.

3.3.2 Results for signal > noise experiments

In DGPs with larger b_{ij} (β_{22}) coefficients — specifically, coefficient values set to those obtained from empirical estimates of inflation models — the signal associated with the x_{it} (x_{22}) variables exceeds the noise, such that the unrestricted model is expected to be more accurate than the restricted model. In this setting, too, our asymptotic results imply the optimal combination forecast should be more accurate than the unrestricted model forecast, on average. However, relative to the accuracy of the unrestricted model forecast, the gains to combination should be smaller than in DGPs with smaller b_{ij} coefficients.

The results for DGPs 1–3 reported in Table 2 confirm these theoretical implications. At the 1-step ahead horizon, the unrestricted model's average MSE is about 5-6 percent lower than the restricted model's MSE in DGP 1 and 3 experiments and roughly 15 percent lower in DGP 2 experiments. At the 4-step ahead horizon, the unrestricted model is more accurate than the restricted by about 12, 28, and 4 percent

for DGPs 1, 2, and 3.

Combination using the known optimal combination weight α_t^* improves accuracy further, more so for DGP 3 (for which the unrestricted forecasting model is largest) than DGPs 1 and 2 and more so for the 4-step ahead horizon than the 1-step horizon. Consider, for example, the forecast sample $P = 1$. For DGP 2, the known optimal combination forecast's MSE ratios are .839 (1-step) and .716 (4-step), compared to the unrestricted forecast's MSE ratios of, respectively, .845 and .723. For DGP 3, the known optimal combination forecast's MSE ratios are .924 (1-step) and .919 (4-step), compared to the unrestricted forecast's MSE ratios of, respectively, .947 and .971. Consistent with our theoretical results, the gains to combination seem to be larger under conditions that likely reduce parameter estimation precision (more variables and residual serial correlation created by the multi-step forecast horizon).

Similarly, the gains to combination (gains relative to the unrestricted model's forecast) rise as the estimation sample gets smaller. Table 3 reports results for the same DGPs used in Table 2, but for the case in which the initial estimation sample is 40 observations instead of 80. With the smaller estimation sample, DGP 2 simulations yield known optimal combination MSE ratios of .882 (1-step) and .807 (4-step), compared to the unrestricted forecast's MSE ratios of, respectively, .908 and .851. For DGP 3, the known optimal combination forecast's MSE ratios are .960 (1-step) and .959 (4-step), compared to the unrestricted forecast's MSE ratios of, respectively, 1.064 and 1.146.

Again, not surprisingly, having to estimate the optimal combination weight tends to slightly reduce the gains to combination. For instance, in Table 2's results for case DGP 2 and $P = 1$, the 4-step ahead MSE ratio for the estimated optimal combination forecast is .723, compared to .716 for the known optimal combination forecast. Using the Stein rule-based adjustment to the optimal combination estimate (based on equation (6)) typically reduces forecast accuracy a bit more (to a MSE ratio of .732 in the same example), but not always — the adjustment often improves forecast accuracy with DGP 3 and a small estimation sample (Table 3).

Imposing simple equal weights in averaging the unrestricted and restricted model forecasts sometimes slightly improves upon the estimated optimal combination but other times reduces accuracy. In Table 2's results for DGPs 1 and 2, the estimated optimal combination is always more accurate than the simple average. For example,

with DGP 2 and the 4-step horizon, the $P = 20$ MSE ratio of the estimated optimal combination forecast is .725, compared to the simple average forecast’s MSE ratio of .767. But for DGP 3, the simple average is often slightly more accurate than the estimated optimal combination. For instance, at the 4-step horizon and with $P = 20$, the optimal combination and simple average forecast MSEs are, respectively, .928 and .919.

As these results suggest, the merits of imposing equal combination weights over estimating weights depend on how far the true optimal weight is from $1/2$ (which depends on the population size and precision of the model coefficients) and the precision of the estimated combination weight. In cases in which the known optimal weight is relatively close to $1/2$ (DGP 3, 1-step forecast, Table 2), the simple average performs quite similarly to the known optimal forecast, and better than the estimated optimal combination. In cases in which the known optimal weight is far from $1/2$ (DGP 2, 1-step forecast, Table 2), the simple average is dominated by the known optimal forecast and, in turn, the estimated optimal combination. Consistent with such reasoning, reducing the initial estimation sample generally improves the accuracy of the simple average forecast relative to the estimated optimal combination. For example, Table 3 shows that, with DGP 2 and the 4-step horizon, the $P = 20$ MSE ratio of the simple average forecast is .789, compared to the estimated optimal combination forecast’s MSE ratio of .775 (in Table 2, the corresponding figures are .767 and .725).

3.3.3 Distributional results

In addition to helping to lower the average forecast MSE, combination of restricted and unrestricted forecasts helps to tighten the distribution of relative accuracy — specifically, the MSE relative to the MSE of the restricted model. The results in Table 4 indicate that combination — especially simple averaging — often increases the probability of equaling or beating the MSE of the restricted model, often by more than it lowers average MSE (note that, to conserve space, the table omits results for DGP 1). For instance, with DGP 2 parameterized such that signal = noise for forecasting 1-step ahead to period $T - P + 2$, the frequency with which the unrestricted model’s MSE is less than or equal to the restricted model’s MSE is 47.2 percent for $P = 20$. The frequency with which the known optimal combination forecast’s MSE is below the restricted model’s MSE is 57.4 percent. Although the

estimated combination does not fare as well (probability of 51.4 percent), a simple average fares even better, beating the MSE of the restricted model in 58.1 percent of the simulations. Note also that, by this distributional metric, using the Stein variant of the combination weight estimate often offers a material advantage over the direct approach to estimating the combination weight. In the same example, the Stein-based combination forecast has a probability of 55.3 percent, compared to the 51.4 percent for the directly estimated combination forecast.

By this probability metric, the simple average (and, to a lesser extent, the optimal combination based on the Stein rule estimate) also fares well in other experiments. Consider, for example, the experiments with the signal > noise version of DGP 3, a forecast horizon of 4 steps, and $P = 20$. In this case, the probability the unrestricted model yields a MSE less than or equal to the restricted model's MSE is 54.3 percent. The probabilities for the estimated optimal combination, Stein-estimated optimal combination, and simple average are, respectively, 62.9, 64.9, and 69.1 percent. Again, averaging, especially simple averaging, greatly improves the probability of beating the accuracy of the restricted model forecast.

4 Empirical Applications

To evaluate the empirical performance of our proposed forecast methods, we consider the widely studied problem of forecasting inflation with Phillips curve models. In particular, we examine forecasts of quarterly core PCE (U.S.) inflation. In light of the potential for the benefits of forecast combination to rise as the number of variables and, in turn, overall parameter estimation imprecision increases, we consider a range of applications, including between one and five predictors of core inflation. In a first application, patterned on analyses in such studies as Stock and Watson (1999, 2003), Orphanides and van Norden (2005), and Clark and McCracken (2006), the unrestricted forecasting model includes lags of inflation and the output gap. In a second application, the unrestricted forecasting model is augmented to include lags of food and energy price inflation, following Gordon's (1998) approach of including supply shock measures in the Phillips curve. In another set of applications, patterned on such studies as Stock and Watson (2002, 2005), and Boivin and Ng (2005), the unrestricted forecast model includes lags of inflation and 1, 2, 3, or 5 common business

cycle factors, estimated as in Stock and Watson (2005). This section proceeds by detailing the data and forecasting models, describing some additional forecast methods included for comparison, and presenting the results.

4.1 Data and model details

Inflation is measured in annualized percentage terms (as 400 times the log change in the price index). The output gap is measured as the log of real GDP less the log of CBO’s estimate of potential GDP. Following Gordon (1998), the food and energy price inflation variable is measured as overall PCE inflation less core PCE inflation. The common factors are estimated with the principal component approach of Stock and Watson (2002, 2005), using a data set of 127 monthly series nearly identical to Stock and Watson’s (2005).¹¹ Following Stock and Watson (2005) and Boivin and Ng (2005), the factors are estimated recursively for each month of the forecast sample, applying the factor estimation algorithm to data through the given month. Quarterly data on factors used in model estimation ending in quarter t are within-quarter averages of monthly factors estimated with data from the beginning of the sample through the last month of quarter t .

Following the basic approach of Stock and Watson (1999, 2003), among others, we treat inflation as having a unit root, and forecast a measure of the direct multi-step change in inflation as a function of lags of the change in quarterly inflation and lags of other variables. In particular, we make y the log difference of the quarterly core PCE price index (scaled by 400 to make y an annualized percentage change); Δy is then the change in quarterly inflation. The predictand is $y_{t+\tau}^{(\tau)} - y_t$, where $y_{t+\tau}^{(\tau)}$ denotes the average annual rate of price change from t to $t + \tau$. The x variables denote the output gap, relative food and energy price inflation, and the set of common factors included in the model (with the number ranging from 1 to 5). The restricted model is autoregressive — the multi-step change in inflation is a function of just lags of the one-period change in inflation. The unrestricted model adds lags of x variables to the set of regressors. In particular, the competing forecasting models take the forms of section 4’s equations (10) and (11). All models include four lags of the change

¹¹Following the specifications of Stock and Watson (2005), we first transformed the data for stationarity, screened for outliers, and standardized the data, and then computed principal components at the monthly frequency.

in inflation (Δy_t). For the output gap and the factors, the models use one lag. For food–energy inflation, the models include two lags.

The forecasting models are estimated with data starting in 1961:Q1. The parameters of the forecasting models are re-estimated with added data as forecasting moves forward through time (that is, our forecasting scheme is the so-called recursive). The forecast sample is 1985:Q1 (1985:Q4 for four-step ahead forecasts) through 2006:Q2. We report results — MSEs — for forecast horizons of one quarter and one year.¹²

4.2 Additional forecast methods

Because our proposed forecast combination methods correspond to a form of shrinkage, for comparison we supplement our results to include not only our proposed methods but also some alternative shrinkage forecasts based on Bayesian methods. Doan, Litterman, and Sims (1984) suggest that conventional Bayesian estimation (specifically, the prior) provides a flexible method for balancing the tradeoff between signal and parameter estimation noise.

Accordingly, one alternative forecast is obtained from the unrestricted forecasting model estimated with generalized ridge regression, which is similar to and under some implementations identical to conventional BVAR estimation. Consistent with the spirit of our proposed combination approaches, which try to limit the effects of sampling noise in the coefficients of the x variables, the ridge estimator pushes the coefficients on the x variables toward zero by imposing informative prior variances on the associated coefficients (the tightness of the prior increases with the number of lags of x included). The ridge estimator allows very large variances on the coefficients of the intercept and lagged inflation terms. In the case of the 1-step ahead model, our generalized ridge estimator is exactly the same as the conventional Bayesian or BVAR estimator of Litterman (1986), except that we use flat priors on the intercept and lagged inflation terms.¹³ We apply the same priors to the 4-step ahead model (based on some experimentation to ensure the prior setting worked well).

We report a second alternative forecast constructed by applying Bayesian model

¹²Because the multi-step forecasts are projections of the average inflation rate from $t + 1$ to $t + \tau$ rather than just the quarterly inflation rate in period $t + \tau$, how forecast accuracy should relate to horizon is unclear. Depending on the DGP, MSEs may rise or fall as the horizon increases.

¹³In the notation of Litterman, we use the following parameter settings in determining the prior variances: $\lambda = .2$ and $\theta = .5$.

averaging (BMA) to the restricted and unrestricted models, following the BMA approach of Fernandez, Ley, and Steel (2001). In particular, we first estimate the models imposing a simple g -prior (but with a flat prior on intercepts), and then average the models based on posterior probabilities calculated as in Fernandez, Ley, and Steel (2001). Following Wright (2003), we set the g -prior coefficient (g_{0j} in the notation of Fernandez, et al., or $1/\phi$ in Wright's notation) at .20.

4.3 Results

In very broad terms, the results in Table 5 seem reasonably reflective of the overall literature on forecasting U.S. inflation in data since the mid-1980s: the variables included in the unrestricted model but not the restricted only sometimes improve forecast accuracy. Across the 12 columns of Table 5 (covering six applications and two forecast horizons), the restricted model's MSE is lower than the unrestricted model's in six cases, sometimes slightly (e.g., 1-year ahead forecasts from the model with five factors) and sometimes dramatically (e.g., 1-year ahead forecasts from the model with the output gap and food-energy inflation). Such a pattern is also consistent with our concept of weak predictability: with a signal-noise ratio that is about 1, such that the restricted and unrestricted models are about equally accurate, we would expect the unrestricted model to beat the restricted about 1/2 of the time (see, e.g., the Monte Carlo results in the top panel of Table 4).

Combining forecasts with our proposed methods significantly improves upon the accuracy of the unrestricted model's forecast, by enough that, in each column, at least one of the average forecasts is more accurate than the restricted model's forecast. For every application and horizon, our estimated optimal combination forecast has a lower MSE than the unrestricted model. For example, in the three factor application (lower block, middle), the optimal combination forecast has a 1-year ahead MSE ratio of .791, while the unrestricted model has a MSE ratio of .879. Consistent with our theoretical results, the advantage of the combination forecast over the unrestricted forecast tends to rise as the number of x variables in the unrestricted model increases (with the increase in the number of variables tending to lower the signal-noise ratio) and as the forecast horizon increases. For example, in the same (three factor) application, the 1-quarter ahead MSE ratios of the unrestricted and optimal combination forecasts are, respectively, .950 and .935 — closer together than

for the 1-year ahead horizon. In the five factor application, the 1-year ahead MSE ratios of the unrestricted and optimal combination forecasts are 1.001 and .834 — farther apart than in the three factor application.

Estimating the optimal combination weight with our proposed Stein rule-based approach yields a consistent, modest improvement in forecast accuracy. In all columns of Table 5, the optimal combination forecast based on the Stein-estimated weight (6) has a lower MSE than does the optimal combination based on the baseline approach (5). In the same three factor application, at the 1-year horizon the optimal combination based on the Stein-estimated weight has a MSE ratio of .781, compared to the directly estimated optimal combination forecast's MSE ratio of .791. With five factors and the 1-year horizon, the Stein-estimated optimal combination's MSE ratio is .807, while the directly estimated optimal combination forecast's MSE ratio is .834.

In most cases, imposing equal weights in combining the restricted and unrestricted model forecasts further improves forecast accuracy, sometimes substantially. As a result, in many cases, the simple average forecast is the best forecast of all considered. For instance, in the output gap and food-energy inflation application, the 1-year ahead MSE ratios of the simple average and Stein-estimated combination forecasts are .871 (the lowest among all forecasts) and 1.020, respectively. In the application with two factors, the 1-year ahead MSE ratios are .854 (again, the lowest among all forecasts) and .866 for, respectively, the simple average and Stein-estimated combination forecasts. In some cases, though, the simple average is only slightly better than or worse than our proposed Stein rule-based approach. For example, in the three factor application, the simple average forecast's 1-year ahead MSE ratio is .782, compared to the Stein-estimated combination forecast's MSE ratio of .781.

In these applications, our proposed combinations clearly dominate Bayesian model averaging and are generally about as good as or better than ridge regression. For example, in the two factor application, the 1-year ahead MSE ratio is .866 for the Stein-estimated combination, .854 for the simple average, and .891 for the ridge regression forecast. In the same application, though, the 1-quarter ahead MSE ratios are virtually identical, at .953, .950, and .950, respectively. In the application with the output gap and food-energy inflation as predictands, the 1-quarter ahead MSE ratios of the Stein-estimated, simple average, and ridge regression forecasts are, respectively, 1.060, .983, and 1.032. However, in all cases, the BMA forecast is less accurate

than the Stein-estimated combination and simple average forecasts. For instance, in the two factor application, the BMA forecast has MSE ratios of .971 and .934 at the 1-quarter and 1-year ahead horizons (compared, e.g., to the Stein-estimated combination MSE ratios of .953 and .866).

5 Conclusion

As reflected in the principle of parsimony, when some variables are truly but weakly related to the variable being forecast, having the additional variables in the model may detract from forecast accuracy, because of parameter estimation error. Focusing on such cases of weak predictability, we show that combining the forecasts of the parsimonious and larger models can improve forecast accuracy. We first derive, theoretically, the optimal combination weight and combination benefit. In the special case in which the coefficients on the variables of interest are of a magnitude that makes the restricted and unrestricted models equally accurate, the MSE-minimizing forecast is a simple, equally-weighted average of the restricted and unrestricted forecasts.

A range of Monte Carlo experiments and empirical examples show our proposed approach of combining forecasts from nested models to be effective in practice. Admittedly, the gains to averaging are often modest or even small. However, the gains are very consistent: in practice, in our results, averaging is very likely to improve on the accuracy of both the restricted and unrestricted model forecasts. Moreover, in practice, most of the benefits can be achieved at low cost, via simple, equal-weight averages. These simple averages typically perform at least as well as more complicated averages. Our paper thus supports the conventional wisdom that simple averages are hard to beat — but, in contrast to most of the combination literature, provides a theoretical and practical basis for applying averaging to nested models.

References

- Atkeson, A. and Ohanian, L. E. (2001). ‘Are Phillips curves useful for forecasting inflation?’, *Quarterly Review*, Federal Reserve Bank of Minneapolis, Vol. 25, pp. 2-11.
- Bates, J. M. and Granger, C. W. J. (1969). ‘The combination of forecasts’, *Operations Research Quarterly*, Vol. 20, pp. 451-468.
- Boivin, J. and Ng, S. (2005). ‘Understanding and comparing factor-based forecasts’, *International Journal of Central Banking*, Vol. 1, pp. 117-151.
- Clark, T. E. and McCracken, M. W. (2005). ‘Evaluating direct multistep forecasts’, *Econometric Reviews*, Vol. 24, pp. 369-404.
- Clark, T. E. and McCracken, M. W. (2006). ‘The predictive content of the output gap for inflation: resolving in-sample and out-of-sample evidence’, *Journal of Money, Credit, and Banking*, Vol. 38, pp. 1127-1148.
- Clark, T. E. and McCracken, M. W. (2006). ‘Combining forecasts from nested models’, Research Working Paper 06-02, Federal Reserve Bank of Kansas City.
- Clements, M. P. and Hendry, D. F. (1998). *Forecasting Economic Time Series*, Cambridge University Press, Cambridge.
- de Jong, R. M. and Davidson, J. (2000). ‘The functional central limit theorem and weak convergence to stochastic integrals I: weakly dependent processes’, *Econometric Theory*, Vol. 16, pp. 621-642.
- Diebold, F. X. (1998). *Elements of Forecasting*, South-Western College Publishing, Cincinnati, OH.
- Doan, T., Litterman, R. and Sims, C. (1984). ‘Forecasting and conditional prediction using realistic prior distributions’, *Econometric Reviews*, Vol. 3, pp. 1-100.
- Elliott, G. and Timmermann, A. (2004). ‘Optimal forecast combinations under general loss functions and forecast error distributions’, *Journal of Econometrics*, Vol. 122, pp. 47-79.
- Fernandez, C., Ley, E. and Steel, M. F. J. (2001). ‘Benchmark priors for Bayesian model averaging’, *Journal of Econometrics*, Vol. 100, pp. 381-427.
- Godfrey, L. G. and Orme, C. D. (2004). ‘Controlling the finite sample significance levels of heteroskedasticity-robust tests of several linear restrictions on regression coefficients’, *Economics Letters*, Vol. 82, pp. 281-287.
- Gordon, R. J. (1998). ‘Foundations of the Goldilocks economy: supply shocks and the time-varying NAIRU’, *Brookings Papers on Economic Activity* (no. 2). pp. 297-346.

- Goyal, A. and Welch, I. (2003). ‘Predicting the equity premium with dividend ratios’, *Management Science*, Vol. 49, pp. 639-654.
- Hansen, B. E. (2008). ‘Averaging estimators for autoregressions with a near unit root’, *Journal of Econometrics*, forthcoming.
- Hendry, D. F. and Clements, M. P. (2004). ‘Pooling of forecasts’, *Econometrics Journal*, Vol. 7, pp. 1-31.
- Jacobson, T. and Karlsson, S. (2004). ‘Finding good predictors for inflation: a Bayesian model averaging approach’, *Journal of Forecasting*, Vol. 23, pp. 479-496.
- Koop, G. and Potter, S. (2004). ‘Forecasting in dynamic factor models using Bayesian model averaging’, *Econometrics Journal*, Vol. 7, pp. 550-565.
- Litterman, R. B. (1986). ‘Forecasting with Bayesian vector autoregressions — five years of experience’, *Journal of Business and Economic Statistics*, Vol. 4, pp. 25-38.
- Orphanides, A. and van Norden, S. (2005). ‘The reliability of inflation forecasts based on output gap estimates in real time’, *Journal of Money, Credit, and Banking*, Vol. 37, pp. 583-601.
- Smith, J. and Wallis, K. F. (2007). ‘A simple explanation of the forecast combination puzzle’, *Oxford Bulletin of Economics and Statistics*, forthcoming.
- Stock, J. H. and Watson, M. W. (1999). ‘Forecasting inflation’, *Journal of Monetary Economics*, Vol. 44, pp. 293-335.
- Stock, J. H. and Watson, M. W. (2002). ‘Macroeconomic forecasting using diffusion indexes’, *Journal of Business and Economic Statistics*, Vol. 20, pp. 147-162.
- Stock, J. H. and Watson, M. W. (2003). ‘Forecasting output and inflation: the role of asset prices’, *Journal of Economic Literature*, Vol. 41, pp. 788-829.
- Stock, J. H. and Watson, M. W. (2005). “An Empirical Comparison of Methods for Forecasting Using Many Predictors,” manuscript, Harvard University.
- Theil, H. (1971). *Principles of Econometrics*, John Wiley Press, New York.
- Wright, J. H. (2003). ‘Forecasting U.S. inflation by Bayesian model averaging’, manuscript, Board of Governors of the Federal Reserve System.

Table 1. Monte Carlo Results from Signal = Noise Experiments: Average MSEs
*(for restricted model, average MSE; for other forecasts,
ratio of average MSE to restricted model's average MSE)*

method/model	DGP 1							
	horizon = 1				horizon = 4			
	$P=1$	$P=20$	$P=40$	$P=80$	$P=1$	$P=20$	$P=40$	$P=80$
restricted	.773	.775	.771	.764	.818	.816	.808	.796
unrestricted	1.004	1.002	1.000	.998	1.029	1.011	1.006	.998
opt. combination: known α_t^*	.995	.994	.994	.993	.995	.986	.985	.982
opt. combination: $\hat{\alpha}_t^*$.999	.998	.997	.996	1.007	.996	.993	.989
opt. combination: Stein $\hat{\alpha}_t^*$.999	.998	.998	.997	1.005	.996	.994	.991
simple average	.995	.994	.994	.993	.992	.984	.983	.981
method/model	DGP 2							
	horizon = 1				horizon = 4			
	$P=1$	$P=20$	$P=40$	$P=80$	$P=1$	$P=20$	$P=40$	$P=80$
restricted	.678	.678	.677	.672	.635	.632	.627	.620
unrestricted	1.009	1.003	.999	.993	1.004	1.004	.996	.983
opt. combination: known α_t^*	.984	.983	.982	.980	.959	.962	.960	.956
opt. combination: $\hat{\alpha}_t^*$.992	.989	.987	.984	.972	.974	.971	.964
opt. combination: Stein $\hat{\alpha}_t^*$.993	.990	.989	.987	.975	.976	.973	.967
simple average	.984	.982	.982	.980	.958	.960	.959	.956
method/model	DGP 3							
	horizon = 1				horizon = 4			
	$P=1$	$P=20$	$P=40$	$P=80$	$P=1$	$P=20$	$P=40$	$P=80$
restricted	.780	.752	.747	.740	.843	.822	.817	.805
unrestricted	1.012	1.009	1.003	.994	1.058	1.050	1.037	1.020
opt. combination: known α_t^*	.974	.974	.973	.970	.972	.973	.971	.967
opt. combination: $\hat{\alpha}_t^*$.983	.982	.980	.976	.993	.991	.987	.980
opt. combination: Stein $\hat{\alpha}_t^*$.985	.983	.981	.978	.987	.985	.983	.978
simple average	.974	.974	.973	.970	.974	.974	.972	.967

Notes:

1. DGPs 1–3 are defined in, respectively, equations (7), (8), and (9). In all experiments, the b_{ij} coefficients are scaled such that the null and alternative models are (in population) expected to be equally accurate in the first forecast period. For DGP 1, $b_{11} = .042$. For DGP 2, $b_{11} = .026, b_{21} = .100, b_{22} = .037$. For DGP 3, $b_{11} = .026, b_{21} = .06, b_{31} = .106, b_{41} = .026, b_{51} = .053$.

2. The forecasting approaches are defined as follows. The *restricted* forecast is obtained from OLS estimates of the model omitting x terms (equation (10)). The *unrestricted* forecast is obtained from OLS estimates of the full model (equation (11)). The *opt. combination: known* α_t^* forecast is computed as $\alpha_t^* \times \text{restricted} + (1 - \alpha_t^*) \times \text{unrestricted}$, with α_t^* computed according to (4), using the known features of the DGP. The *opt. combination: $\hat{\alpha}_t^*$* forecast is $\hat{\alpha}_t^* \times \text{restricted} + (1 - \hat{\alpha}_t^*) \times \text{unrestricted}$, with $\hat{\alpha}_t^*$ computed according to (8), using moments estimated from the data. The *opt. combination: Stein $\hat{\alpha}_t^*$* forecast is $\hat{\alpha}_t^* \times \text{restricted} + (1 - \hat{\alpha}_t^*) \times \text{unrestricted}$, with $\hat{\alpha}_t^*$ computed according to (9). Finally, the *simple average* forecast is $.5 \times \text{restricted} + .5 \times \text{unrestricted}$.

3. P defines the number of observations in the forecast sample. The size of the sample used to generate the first (in time) forecast at horizon τ is $80 - \tau + 1$ (the estimation sample expands as forecasting moves forward in time).

4. The table entries are based on averages of forecast MSEs across 10,000 Monte Carlo simulations.

Table 2. Monte Carlo Results from Signal > Noise Experiments: Average MSEs
*(for restricted model, average MSE; for other forecasts,
ratio of average MSE to restricted model's average MSE)*

method/model	DGP 1							
	horizon = 1				horizon = 4			
	$P=1$	$P=20$	$P=40$	$P=80$	$P=1$	$P=20$	$P=40$	$P=80$
restricted	.813	.813	.809	.802	.958	.955	.946	.932
unrestricted	.955	.954	.953	.950	.898	.882	.878	.871
opt. combination: known α_t^*	.952	.952	.951	.949	.889	.875	.872	.867
opt. combination: $\hat{\alpha}_t^*$.957	.955	.954	.952	.895	.881	.878	.872
opt. combination: Stein $\hat{\alpha}_t^*$.960	.958	.957	.954	.903	.888	.884	.877
simple average	.959	.959	.958	.958	.896	.890	.889	.887
method/model	DGP 2							
	horizon = 1				horizon = 4			
	$P=1$	$P=20$	$P=40$	$P=80$	$P=1$	$P=20$	$P=40$	$P=80$
restricted	.811	.805	.803	.798	.967	.948	.940	.929
unrestricted	.845	.845	.841	.837	.723	.729	.724	.714
opt. combination: known α_t^*	.839	.840	.837	.834	.716	.721	.718	.710
opt. combination: $\hat{\alpha}_t^*$.843	.843	.840	.836	.723	.725	.722	.713
opt. combination: Stein $\hat{\alpha}_t^*$.847	.845	.841	.837	.732	.732	.728	.718
simple average	.865	.866	.866	.865	.765	.767	.767	.764
method/model	DGP 3							
	horizon = 1				horizon = 4			
	$P=1$	$P=20$	$P=40$	$P=80$	$P=1$	$P=20$	$P=40$	$P=80$
restricted	.834	.803	.798	.790	.968	.942	.934	.921
unrestricted	.947	.944	.939	.931	.971	.966	.956	.940
opt. combination: known α_t^*	.924	.924	.922	.918	.919	.920	.917	.910
opt. combination: $\hat{\alpha}_t^*$.930	.928	.926	.921	.929	.928	.923	.915
opt. combination: Stein $\hat{\alpha}_t^*$.936	.932	.929	.924	.935	.932	.928	.920
simple average	.928	.927	.927	.925	.918	.919	.917	.913

Notes:

1. DGPs 1–3 are defined in, respectively, equations (7), (8), and (9). For DGP 1, $b_{11} = .042$. For DGP 2, $b_{11} = .07, b_{21} = .27, b_{22} = .10$. For DGP 3, $b_{11} = .04, b_{21} = .09, b_{31} = .16, b_{41} = .04, b_{51} = .08$.
2. See the notes to Table 1.

Table 3. Signal > Noise Experiments with Small Estimation Sample: Average MSEs
*(for restricted model, average MSE; for other forecasts,
ratio of average MSE to restricted model's average MSE)*

method/model	DGP 1							
	horizon = 1				horizon = 4			
	$P=1$	$P=20$	$P=40$	$P=80$	$P=1$	$P=20$	$P=40$	$P=80$
restricted	.854	.853	.839	.823	1.061	1.023	.998	.966
unrestricted	.988	.970	.964	.958	.970	.938	.918	.900
opt. combination: known α_t^*	.971	.961	.957	.954	.925	.909	.897	.886
opt. combination: $\hat{\alpha}_t^*$.978	.967	.962	.958	.924	.913	.902	.891
opt. combination: Stein $\hat{\alpha}_t^*$.980	.971	.967	.962	.925	.919	.909	.898
simple average	.968	.962	.961	.959	.908	.902	.897	.894
method/model	DGP 2							
	horizon = 1				horizon = 4			
	$P=1$	$P=20$	$P=40$	$P=80$	$P=1$	$P=20$	$P=40$	$P=80$
restricted	.876	.852	.837	.820	1.071	1.024	.998	.970
unrestricted	.908	.882	.868	.855	.851	.801	.775	.749
opt. combination: known α_t^*	.882	.865	.856	.847	.807	.773	.755	.736
opt. combination: $\hat{\alpha}_t^*$.888	.870	.860	.850	.804	.775	.758	.741
opt. combination: Stein $\hat{\alpha}_t^*$.896	.878	.866	.854	.819	.791	.773	.751
simple average	.886	.877	.873	.870	.807	.789	.782	.775
method/model	DGP 3							
	horizon = 1				horizon = 4			
	$P=1$	$P=20$	$P=40$	$P=80$	$P=1$	$P=20$	$P=40$	$P=80$
restricted	.841	.837	.827	.812	1.022	1.009	.988	.960
unrestricted	1.064	1.019	.993	.967	1.146	1.087	1.047	1.002
opt. combination: known α_t^*	.960	.950	.941	.932	.959	.952	.943	.929
opt. combination: $\hat{\alpha}_t^*$.980	.963	.951	.939	.994	.976	.961	.942
opt. combination: Stein $\hat{\alpha}_t^*$.972	.962	.953	.942	.968	.962	.953	.940
simple average	.957	.947	.940	.934	.964	.951	.941	.929

Notes:

1. DGPs 1–3 are defined in, respectively, equations (7), (8), and (9). For DGP 1, $b_{11} = .042$. For DGP 2, $b_{11} = .07, b_{21} = .27, b_{22} = .10$. For DGP 3, $b_{11} = .04, b_{21} = .09, b_{31} = .16, b_{41} = .04, b_{51} = .08$.
2. P defines the number of observations in the forecast sample. The size of the sample used to generate the first (in time) forecast at horizon τ is $40 - \tau + 1$ (rather than $80 - \tau + 1$ as in the baseline experiments).
3. See the notes to Table 1.

**Table 4: Monte Carlo Probabilities of Equaling
or Beating Restricted Model’s MSE, DGPs 2 and 3**

method/model	DGP 2: signal = noise							
	horizon = 1				horizon = 4			
	$P=1$	$P=20$	$P=40$	$P=80$	$P=1$	$P=20$	$P=40$	$P=80$
unrestricted	.501	.472	.483	.524	.509	.495	.493	.525
opt. combination: known α_t^*	.521	.574	.627	.698	.535	.576	.600	.660
opt. combination: $\hat{\alpha}_t^*$.515	.514	.547	.613	.530	.538	.554	.605
opt. combination: Stein $\hat{\alpha}_t^*$.642	.553	.554	.592	.646	.597	.579	.601
simple average	.521	.581	.639	.727	.539	.593	.628	.706
method/model	DGP 2: signal > noise							
	horizon = 1				horizon = 4			
	$P=1$	$P=20$	$P=40$	$P=80$	$P=1$	$P=20$	$P=40$	$P=80$
unrestricted	.557	.803	.901	.977	.589	.785	.869	.951
opt. combination: known α_t^*	.567	.834	.926	.986	.600	.810	.893	.963
opt. combination: $\hat{\alpha}_t^*$.570	.843	.935	.989	.609	.836	.916	.975
opt. combination: Stein $\hat{\alpha}_t^*$.574	.849	.939	.990	.618	.844	.921	.977
simple average	.598	.921	.980	.999	.635	.900	.965	.995
method/model	DGP 3: signal = noise							
	horizon = 1				horizon = 4			
	$P=1$	$P=20$	$P=40$	$P=80$	$P=1$	$P=20$	$P=40$	$P=80$
unrestricted	.497	.461	.473	.519	.481	.424	.412	.417
opt. combination: known α_t^*	.524	.609	.652	.736	.521	.564	.590	.642
opt. combination: $\hat{\alpha}_t^*$.517	.549	.584	.670	.505	.506	.513	.555
opt. combination: Stein $\hat{\alpha}_t^*$.608	.564	.583	.664	.614	.554	.540	.562
simple average	.524	.616	.671	.770	.517	.556	.587	.653
method/model	DGP 3: signal > noise							
	horizon = 1				horizon = 4			
	$P=1$	$P=20$	$P=40$	$P=80$	$P=1$	$P=20$	$P=40$	$P=80$
unrestricted	.521	.621	.684	.796	.512	.543	.571	.648
opt. combination: known α_t^*	.541	.715	.794	.899	.540	.644	.696	.786
opt. combination: $\hat{\alpha}_t^*$.539	.702	.778	.890	.537	.629	.677	.773
opt. combination: Stein $\hat{\alpha}_t^*$.568	.710	.791	.900	.587	.649	.688	.787
simple average	.554	.779	.867	.955	.552	.691	.758	.861

Notes:

1. The table entries are frequencies (percentages of 10,000 Monte Carlo simulations) with which each forecast approach yields a forecast MSE less than or equal to the restricted model’s MSE.
2. See the notes to Tables 1 and 2.

Table 5. Application Results: 1985-2006 Forecasts of Core PCE Inflation
*(for restricted model, average MSE; for other forecasts,
ratio of MSE to restricted model's MSE)*

method/model	output gap		output gap & food-energy inflation		1 factor	
	1Q	1Y	1Q	1Y	1Q	1Y
restricted	.632	.516	.632	.516	.632	.516
unrestricted	.980	1.044	1.150	1.380	.979	1.034
opt. combination: $\hat{\alpha}_t^*$.976	.990	1.073	1.081	.977	.986
opt. combination: Stein $\hat{\alpha}_t^*$.976	.984	1.060	1.020	.977	.978
simple average	.973	.906	.983	.871	.977	.950
Ridge regression	.978	1.018	1.032	1.135	.976	.976
BMA	.995	1.006	1.085	1.155	.999	1.024
method/model	2 factors		3 factors		5 factors	
	1Q	1Y	1Q	1Y	1Q	1Y
restricted	.632	.516	.632	.516	.632	.516
unrestricted	.965	.971	.950	.879	1.136	1.001
opt. combination: $\hat{\alpha}_t^*$.954	.879	.935	.791	1.040	.834
opt. combination: Stein $\hat{\alpha}_t^*$.953	.866	.934	.781	1.021	.807
simple average	.950	.854	.936	.782	.963	.794
Ridge regression	.950	.891	.933	.807	.955	.815
BMA	.971	.934	.954	.846	1.065	.914

Notes:

1. The forecasting models take the forms given in equations (10) and (11). In the first application, the unrestricted model includes just one lag of the output gap, defined as the log ratio of actual GDP to the CBO's estimate of potential GDP. In the second application, the unrestricted model includes one lag of the output gap and two lags of relative food and energy price inflation, calculated as overall PCE inflation less core PCE inflation. In the remaining applications, the unrestricted model includes one lag of common business cycle factors — with the number of factors varying from 1 to 5 across applications — estimated as in Stock and Watson (2005).
2. The first six forecast approaches are defined in the notes to Table 1. The *BMA* forecast is a Bayesian average of the forecasts from the restricted and unrestricted models, implemented with the averaging approach recommended by Fernandez, Ley, and Steel (2001), with the difference that these results are based on a *g*-prior coefficient setting of 1/5. The *Ridge regression* forecast is obtained from a generalized ridge estimator which shrinks the β_{22} coefficients (but not the other coefficients) of the unrestricted model toward 0 based on conventional Minnesota prior settings described in section 4.2.