

2.08.09 Ultraviolet B Modulates Stress Responses of *Lemna minor* to Gamma Radiation at Multiple Levels of Biological Organization

L. Xie, NIVA - Norwegian Institute for Water Research / Section of Ecotoxicology and Risk Assessment; Y. Song, Norwegian Institute for Water Research (NIVA) / Section for Ecotoxicology and Risk Assessment; K. Petersen, NIVA - Norwegian Institute for Water Research / Section of Ecotoxicology and Risk Assessment; K. Solhaug, Norwegian University of Life Sciences (NMBU) / Faculty of Environmental Sciences and Natural Resource Management; O. Lind, Norwegian University of Life Sciences, Center of Excellence in Environmental Radioactivity (CERAD) / Faculty of Environmental Sciences and Natural Resource Management; D.A. Brede, Norwegian University of Life Sciences / Centre for Environmental Radioactivity; B. Salbu, Norwegian University of Life Sciences (NMBU) / Centre for Environmental Radioactivity (CERAD); K. Tollefsen, NIVA - Norwegian Institute for Water Research / Section of Ecotoxicology and Risk Assessment. Climate change, such as global warming, has been shown to have important implications for future stratospheric ozone depletion, which determines the intensity of UV radiation at the Earth's surface. In the aquatic ecosystem, UV radiation and ionizing radiation may co-occur and pose cumulative hazards to aquatic organisms such as macrophytes. However, the combined effects and underlying mechanisms of different types of radiation in aquatic plants remain poorly understood. The present study aims to provide in-depth understanding of how chronic (7 days) exposure to a single irradiance (0.5 W m^{-2}) of ultraviolet B (UVB) radiation modulates gamma (γ) radiation ($14.9, 19.5, 43.6 \text{ mGy h}^{-1}$) induced stress responses in the aquatic macrophyte *Lemna minor*. In contrast to a majority of combined effect studies that focus on either molecular or apical endpoints, the present study used a suite of bioassays to quantify stress responses at multiple levels of biological organization. The combined effects (additivity, synergism, antagonism) of the stressors were determined using a combination of two-way analysis of variance (2W-ANOVA) and a modified Independent Action (IA) model. The single and combined effects of the stressors, and the potential causality between toxicological responses were further visualized by assembly of putative toxicity pathways. The results show that exposure to γ -radiation or UVB alone commonly induced oxidative stress and programmed cell death (PCD), and impaired oxidative phosphorylation (OXPHOS) and photosystem II (PSII) in *L. minor*. γ -radiation also activated antioxidant responses, DNA damage repair and chlorophyll metabolism, and inhibited growth at high dose rates (20 mGy h^{-1}). Additive effects of the two stressors were predominantly identified for antioxidant gene expression, energy quenching in PSII, frond size and frond weight for all dose rates, whereas antagonistic effects were observed for DNA damage, PCD, oxidative stress, OXPHOS and chlorophyll metabolism. Synergistic inhibition was observed for reproduction (fronds number) for all dose rates. The present study provides substantial mechanistic knowledge, quantitative understanding and novel analytical strategies to decipher combined effects at different levels of biological organization and is anticipated to facilitate future cumulative hazard and risk assessment of multiple stressors.