

CLASS: A

PROTOCOL(S) USED IN: Seizures

PHARMACOLOGY AND ACTIONS:

Non-narcotic analgesic and antipyretic

INDICATIONS:

- A. Reduction of fever associated with febrile seizures in the pediatric patient.

CONTRAINDICATIONS:

- A. DO NOT use with any other products that contain acetaminophen

SIDE EFFECTS AND NOTES:

- A. May be administered via rectal suppository (same dose) if patient is vomiting, the patient's gag reflex is absent or in question or the patient is not alert.
- B. Can be given up to q 4 hours.
- C. Do not exceed 5 doses a day.

DOSING CHART:

Weight	Age	80mg Tablets	160mg/tsp Elixir	80mg/0.8ml Drops	Mg
6-11 lbs	0-3 mo	---	¼ tsp	.4ml	40mg
12-17lbs	4-11mo	---	½ tsp	.8ml	80mg
18-23lbs	11-23mo	1 ½ tab	¾ tsp	1.2ml	120mg
24-35lbs	2-3yrs	2 tabs	1 tsp	1.6ml	160mg
36-47lbs	4-5yrs	3 tabs	1 ½ tsp	2.4ml	240mg
48-59lbs	6-8yrs	4 tabs	2 tsp	3.2ml	320mg
60-71lbs	9-10yrs	5 tabs	2 ½ tsp	4.0ml	400mg
72-95lbs	11-12yrs	6 tabs	3 tsp	4.8ml	480mg