
2021 Sec 4 Science Chemistry Prelim - Hillgrove

2

21 In which conversion do H₂O molecules lose speed?



- Α ice → water
- В ice → steam
- C steam → water
- D water → steam
- 22 A student was asked to carry out the following steps in an experiment:

Step I : Hold about 200 cm3 of solution X.

: Transfer 25.0 cm³ of solution X into a conical flask. Step II

Step III : Add 34.60 cm³ of aqueous hydrochloric acid into the conical flask.

Which row lists the most suitable piece of apparatus for each of the above three steps?


	Step I	Step II	Step III
Α	beaker	pipette	burette
В	beaker	burette	pipette
С	conical flask	pipette	measuring cylinder
D	measuring cylinder	burette	measuring cylinder

23 Methanol boils at 65°C and water boils at 100°C. Methanol and water are completely miscible with each other.

Which method is used to separate a mixture of these two liquids?


- Α evaporation
- В filtration
- C fractional distillation
- D paper chromatography

24 The chromatogram below shows three drinks, Fizzy, Juicy and Sparkly, analyzed for any banned colourings.




Which of the drinks contain banned colourings?

- Α Fizzy and Juicy
- В Fizzy and Sparkly
- Juicy and Sparkly C
- D Sparkly only
- 25 The diagram below shows the particles of hexane at atmospheric pressure but at two different temperatures. Hexane melts at -95°C and boils at 70°C.







What could be the values of X and Y?

	X	Υ
A	-100	Ö
В	50	10
С	0	100
D	10	– 50

26 An element Y has two isotopes, ²³⁸Y and ²³⁵Y.

How does ²³⁸Y differ from ²³⁵Y.


- A It has 3 more protons.
- B It has 3 more neutrons.
- C It has 3 more protons and 3 more electrons.
- **D** It has 3 more neutrons and 3 more electrons.
- 27 Which ions are present in Cu₂SO₄?
 - A Cu²⁺, S⁶⁺ and O²⁻
 - B Cu²⁺ and SO₄²⁻
 - C Cu⁺ and SO₄²⁻
 - D Cu²⁺ and SO₄
- 28 Why does molten sodium chloride conduct electricity?
 - A Electrons in the sodium chloride are free to move.
 - B Sodium ions are strongly attracted to the chloride ions.
 - C The sodium ions and the chloride ions are free to move.
 - **D** Electrons in the valence shell of sodium atoms are transferred to chlorine atoms.
- 29 20 cm³ of ethene is reacted with 30 cm³ of hydrogen to form ethane gas. The equation for the reaction is shown.

$$C_2H_4(g) + H_2(g) \rightarrow C_2H_6(g)$$

What is the volume of each of the gases at the end of the reaction? (all volumes measured at r.t.p)

	volume of C ₂ H ₄ / cm ³	volume of H ₂ / cm ³	volume of C ₂ H ₆ / cm ³
Α	0	0	50
В	0	10	20
С	10	0	10
D	0	10	50

30 In which test-tube is the iron nail most likely to rust?



- 31 Which statement is not true about an alloy?
 - An alloy is a mixture. Α
 - An alloy may contain a non-metal. В
 - The chemical properties of an alloy are similar to the elements that form it. C
 - The physical properties of an alloy are similar to the elements that form it. D
- 32 Which reaction is a redox reaction?

B Ag⁺ + C
$$l^ \rightarrow$$
 AgC l

c
$$2SO_2 + O_2 \rightarrow 2SO_3$$


D MgO +
$$2HCl \rightarrow MgCl_2 + H_2O$$

33 An oxide of element V, reacts with both dilute hydrochloric acid and sodium hydroxide solution to form salt and water.

Which row describes V and its oxide?

	V	V oxide
Α	metal	basic
В	metal	amphoteric
С	non-metal	amphoteric
D	non-metal	acidic


34 The diagram shows a reaction scheme for solution X.



What is(are) the anion(s) in solution X?

- A sulfate
- **B** carbonate
- C carbonate and sulfate
- D carbonate and chloride
- **35** Aqueous sodium hydroxide and dilute hydrochloric acid are reacted together. The reaction is exothermic. Both the solutions are initially at room temperature.

Which graph shows how the temperature changes when aqueous sodium hydroxide is added to dilute hydrochloric acid until the alkali is present in excess?



36 Metals W, X, Y and Z are placed in salt solutions as shown in the table.

		result of placing n	netal in solution of			
metals	salt of W	salt of X	salt of Y	salt of Z		
W	no reaction	X displaced	Y displaced	no reaction		
Х	no reaction	no reaction	no reaction	no reaction		
Υ	no reaction	X displaced	no reaction	no reaction		
Z	W displaced	X displaced	Y displaced	no reaction		

What is the order of reactivity of the metals from most reactive to least reactive?

 $X \rightarrow Y \rightarrow W \rightarrow Z$ Α

 $Y \rightarrow W \rightarrow Z \rightarrow X$ В

 $Z \rightarrow W \rightarrow Y \rightarrow X$ C

 $Z \rightarrow Y \rightarrow X \rightarrow W$ D

37 Which substance represents a metal?

	state at room temperature	melting and boiling point	conduction of heat and electricity	
Α	liquid	low	non-malleable	poor
В	solid	high	malleable	good
С	solid	high	malleable	poor
D	solid	low	non-malleable	good

38 A student tests four solutions with Universal Indicator.

solution	P	Q	R	S
colour with universal indicator	purple	green	red	yellow

What are the pH values of the four solutions?

	Р	Q	R	S
Α	2	9	4	7
В	4	7	8	9
С	9	5	12	2
D	12	7	2	5

39 Part of the Periodic Table is shown.


Period			Group					,,	
	1	11		Ш	I۷	٧	۷I	VII	0
1							Ŷ.		
2	٧	W						X	
3	Y							Z	

The letters are not the symbols of the elements.

Which statement is correct?

- V is more reactive than Y.
- W has more metallic character than V. В
- Y has a higher melting point than V. C
- Z is less reactive than X. D
- 40 Some crystals of magnesium carbonate were added to an excess of sulfuric acid at room temperature. The volume of carbon dioxide produced was measured over a period of time. The results are shown in graph Y.

The experiment was repeated and graph **Z** was obtained.



Which change was used to obtain the results shown in graph Z?

- Using a lower temperature. Α
- Half the mass of magnesium carbonate was used. В
- Larger crystals of magnesium carbonate were used. C
- Acid of the same volume and half the original concentration was used. D

9

Data Sheet Colours of Some Common Metal Hydroxides

calcium hydroxide	white
copper(II) hydroxide	light-blue
iron(II) hydroxide	green
iron(III) hydroxide	red-brown
lead(II) hydroxide	white
zinc hydroxide	white

The Periodic Table of Elements

						ar.	I	Trancium	Ţ	87	133	Cacsium	ያ ጸ	8	rubidium	공	37	39	volatesium	ス :		23		<u></u> =	-	7		<u>.</u> ω				-
		aven made	ortinoide.			lanthanoids	ı	Mundey	<u> </u>	88	137		စ္က	88	strontium	र्फ	<u>چ</u>	40	calcium	က္က	20	24	magnesium B	<u> </u>	3,	o y		4 9				=
		•	74			ist.			actinoids	-			3/ - /1	88	ythurn	≺	39	45	scandium	S :	21	•			•							
1	actinium	A (89	139	ត	57	-	Numeriordium	7.7	104	17B		ĘS	9	zircanium	2′	46	48	titanium	= 1	22				15Jan	- Alati	arc	proton				
232	thorium	≓ 8	8	140 140	ဂ္ဂ	85	•						d 3				- 1			< !	1				Courte acomp mass		atomic symbol	proton (atomic) number	Key	•		
231	protactinium	ָּטָ	2	141	P	59	1	Sosporgum	&	106	184	lungston	€ ≥	98	molybdenum	₹	4 2	53	chromium	ល្អ	24				111000		2	nedmin				
238		- f	8	144	ď	60							P 3		technetium	급:	43	ঞ	manganese	<u> </u>	X				-				,			
1		<u>z</u> 8	3	T THE	: 3	2		HUISEE	당	Ē	190		2 %	_						Ţ :	٦								_	hydrogen	I -	
1 Million	olutonium	P 9	2	150	Sm	62	1	melinorium	8	192	j.	77	ន	modium	 공	45	59	SO S	3 :	27											
		A	9	152	旜	63	,	deamstedium	8	110	195	nlafin	₽ 78	106	palladium	D ;	5	ස	nickel	Ζ. \$	8											
1	Curion	<u> </u>	5	gadolmlum	ត្ត	64	1						79																			
1	berkekun	P 4	27	159 159	ฮ	8	,	copomicium	.Ω	112	201	F. 1	E 8	112	cadmium	δ:	48	g,	;	7 6	3											
- Capacitation	californium	2 S	Q.	03300	₽	86		_			204	* -	<u> </u>	115	ndium.	5 ;	49	70		<u>ء</u>	ייי	27)	2 5	=	Doron	0	· Ct				
- CHANGE	cinsleinium	TI S	8	165 765	占	67	1	florovium	. 77	114	207	[]	₽ 83 83	119	5 9	s :	50	73	germanium	9	3	28	2	2 =	_	Carbon	ဂ	a				~
1 6	formi	3 8	3	erbum	щ	68					209	<u> </u>	<u></u> 83	123	antimony	<u>ક</u> ક	51	75	in early	<u> </u>	1	31	7	J 7	,	nifogen	z	7				<
- Interpretation	mendelmin	5 5		160 m	ı M	68	ŀ	livermorium	- -	116	1	7	3 2	128	tellurium	al i	3	79	selenium	វូ វ	2	ខ	 - C	, <u>a</u>	5	oxygen	O	œ				S
1 Indicated	nobelian	<u> </u>	3	ytterbium 173	4	70					1	2	≥ 85	127	odine •	– 8	3	8		₽ {	n N	35.5	ć	? =		nuonna	7	90				≦
1	i i	7 5	103	175	Ε	71		•	•		ı	1 2	0 8	131	xenon	× 4	2	22	Krypion	Ş (3	45	2	2 2	2	7805	Z	ō	4	limital	품 ~	0

HGV Sec 4E/5NA Science (Chemistry) Prelim Paper 1 2021

3

Section A (45 marks)

Answer all the questions in the spaces provided.

1	Iron is	extracted	from	iron	ore	in the	Blast	Furnace.
---	---------	-----------	------	------	-----	--------	-------	----------

The equations A, B, C, D and E show some reactions that happen in the Blast Furnace.

- C + O₂ [] CO₂ Α
- CO₂ + C [2CO В
- C Fe₂O₃ + 3CO [] 2Fe + 3CO₂
- D CaCO₃ CaO + CO₂
- Ε SiO₂ + CaO [] CaSiO₃

(b)

(c)

Use the letters A, B, C, D and E to answer the following equations. You may use each letter once, more than once or not at all.

- Which equation shows combustion? (a)
- [1]
- [1]
- Which equation shows a reaction between an acidic compound and a base?
- [1]
- (d) Which equation shows the formation of a toxic gas?

Which equation shows thermal decomposition?

- [1]
- Which equation shows a redox reaction? (e)
 - [1]

2	Carbon disulfide is a simple covalent compound used in manufacturing polymers and
	fibres.

(a) Draw a 'dot-and-cross' diagram to show the bonding in carbon disulfide.

Show the valence shell only.

[2]

(b) Using your understanding of bonding and structure, which of these statements would you predict to be true or false?

Put a tick (II) in one box in each row.

	True	False
Carbon disulfide has a low boiling point.		
Carbon disulfide is very soluble in water.		
Carbon disulfide has good electrical conductivity when molten.		
Carbon disulfide is a crystalline solid at room temperature.		

[2]

(c) (i) Sulfur reacts with magnesium to form an ionic compound called magnesium sulfide. Draw a 'dot-and-cross' diagrams to show the arrangement of outer shell electrons in its ions and its charges.

magnesium ion

sulfide ion

[2]

(ii)	Explain in terms of bonding, why magnesium sulfide has a high melting and boiling point.				
		[2]			

The rate of reaction between dilute hydrochloric acid and sodium thiosulfate solution 3 can be investigated using a cross drawn on a piece of paper as shown in Figure 3.1.




Figure 3.1

As the reaction progresses, it becomes more difficult to see the cross through the solution.

(a) Look at the equation for the reaction.

 $Na_2S_2O_3(aq) + 2HCl(aq) \square 2NaCl(aq) + S(s) + SO_2(g) + H_2O(l)$

Explain why it becomes more difficult to see the cross as the reaction progresses.

......[2]

(b) Table 3.1 shows the results of an experiment to investigate the rate of reaction using different concentrations of sodium thiosulfate solutions.

A student measured the time from when the acid was added until the cross can no longer be seen.


Table 3.1

Concentration of Na ₂ S ₂ O ₃ / mol/dm ³	Time taken until cross cannot be seen / s
1.0	8
0.8	10
0.4	20
0.2	39

(i)	sodium thiosulfate solution?	
		[1]
(i	Use ideas about collisions between particles to explain the trend given in part (b)(i).	
i)	,	
		[2]
(i i	State two variables that needed to be kept constant in order to make it a fair experiment.	
i)		• • •
		[1]

- Using the results obtained in Table 3.1, plot a graph of concentration of sodium thiosulfate against time taken until the cross cannot be seen.
- Draw a best fit curve through the points.

Concentration of Na₂S₂O₃ / mol/dm³



The experiment was repeated with another concentration of sodium thiosulfate.

The cross could not be seen after 14 s. From your graph in part (iii), suggest the concentration of sodium thiosulfate that was used.

Concentration of sodium thiosulfate: mol/dm3 [1]

Cars have catalytic converters fitted to reduce the problems caused by some of the exhaust gases. Figure 4.1 shows some of the gases that enter and leave a catalytic converter.




Figure 4.1

(a) Carbon monoxide and nitrogen monoxide react in the catalytic converter as shown by the equation below.

Complete Table 4.1 to show the oxidation states of carbon and nitrogen in the reactants and products of the reaction.

Circle the keywords to reach the correct conclusion of the oxidation and reduction of carbon and nitrogen in this reaction.

Table 4.1

	Oxidation state in reactants	Oxidation state in products	Conclusion
carbon	+2		Carbon in carbon monoxide is oxidised / reduced as its oxidation state increased / decreased.
nitrogen		0	Nitrogen in nitrogen monoxide is oxidised / reduced as its oxidation state increased / decreased.

[4]

Briefly describe the harmful effects of nitrogen oxides and carbon monoxide.	
nitrogen oxides :	
carbon monoxide :	
	[2]
Petrol contains mainly octane, C ₈ H ₁₈ which is the main pollutant of unburnt hydrocarbons in exhaust gases of cars. Octane is converted to carbon dioxide and water in the catalytic converters as well as shown by the equation below.	
2C ₈ H ₁₈ + 25O ₂ [] 16CO ₂ + 18H ₂ O	
(Calculate the relative molecular mass of octane. i [Relative atomic masses, A _r : C, 12 ; H, 1])	
M _r of octane =	[1]
 (132g of octane is found in the exhaust gases of an old car. Calculate the number of moles of octane. i) 	
number of moles of octane =	[1]
(Calculate the number of moles of oxygen needed to react with 132g of i octane. i i)	
number of moles of oxygen =	[1]
	carbon monoxide: Carbon monoxide: Petrol contains mainly octane, C ₈ H ₁₈ which is the main pollutant of unburnt hydrocarbons in exhaust gases of cars. Octane is converted to carbon dioxide and water in the catalytic converters as well as shown by the equation below. 2C ₈ H ₁₈ + 25O ₂ 16CO ₂ + 18H ₂ O Calculate the relative molecular mass of octane. [Relative atomic masses, A _r : C, 12; H, 1] M _r of octane =

5 The flow chart in Figure 5.1 shows some reactions of dilute sulfuric acid. Five unknown substances V, W, X, Y and Z are shown in the chart.




Figure 5.1


(a) Suggest the identities of substances V, W, X, Y and Z


i	V		
) (i	w		
) (i	X		
i) (i	Y	••••••	
) (\)	z		[5]
	Vrite an Iowchart	ionic equation with state symbols for any one of the reaction in the	
			[2]

HGV Sec 4E/5NA Science (Chemistry) Prelim Paper 3 2021

(b)

Figure 6.1 shows five boxes describing two solids, two liquids and a gas. (a)





placing a tick (II) in the appropriate column in the table below.

substances	element	compound	mixture
F			
G			
Н			

12

l		
J		[5]

(b)	Describe fully two differences between a compound and a mixture.	
	difference 1 :	
	difference 2:	
		[2

Section B (20 marks)

Answer any two questions in this section. Write your answers in the spaces provided.

Sodi	Sodium has a proton number 11.		
(a)	The Periodic Table lists the elements in groups and periods. An inspection of the electronic structure or configuration of an element's atom can give its group and period. Explain using sodium as an example.		
		[3]	
(b)	Sodium and potassium have similar chemical reactions. Describe two of these similar reactions by referring to their reactants and the type of products they form.		
	Write a chemical equation for one of the reactions using any one of the two metals. Include state symbols.		
		[4]	

(c) Sodium and neon are from two different groups in the Periodic Table and thus have very different chemical reactivity. State the names of the groups they belong to and use their electronic structures to explain the difference in

HGV Sec 4E/5NA Science (Chemistry) Prelim Paper 3 2021

7

reactivity.

			••••••		••••••	
			•••••••		·····	
		*****			• • • • • • • • • • • • • • • • • • • •	[3]
8			ms of chlorine are ide but they can have diffe	entical. All chlorine ato erent masses.	ms have the same ch	emical
	(a)	State	the name given to diff	ferent atoms of the sam	ne element.	
		•••••		•••••		[1]
	(b)		in why different atoms chemical properties?	s of chlorine can have o	lifferent masses but ha	ave the
						•••••
					•••••	
				••••••••••••••••		[2]
	(c)		ifferent masses of the re atomic mass of chlo	chlorine atoms are ave prine which is 35.5.	eraged out to arrive at	the
		(i)	Define the term relati	ve atomic mass.		

						[1]
		(ii)		orine-37 are the two ma eir percentage abundar	•	e. The
				chlorine-35	chlorine-37	
			percentage	75	25	

By using these values, show your working to calculate the relative atomic mass of chlorine.

HGV Sec 4E/5NA Science (Chemistry) Prelim Paper 3 2021

abundance / %

[1]

(d) Draw the electronic structure of an atom of chlorine—35. Show in your structure the number of protons and neutrons as well.

[3]

(e) Chlorine reacts with carbon monoxide to produce phosgene gas with the formula COCl₂.

$$Cl_2(g) + CO(g) \square COCl_2(g)$$

Calculate volume of phosgene than can be produced if 142g of chlorine was used. [Relative atomic masses, A_r : C, 12; O, 16; Cl, 35.5]

[2]

9	(a)	Pure crystals of sodium sulfate can be made from a solution of sodium sulfate.	
		Describe the process of preparing this solution using a solution of sodium hydroxide and an acid.	
			[6]
	(b)	Calculate the mass of sodium hydroxide contained in 250 cm ³ of a solution of sodium hydroxide of concentration of 2.0 mol/dm ³ .	
		[Relative atomic masses, A _r : Na, 23; O, 16; H, 1]	
			[2]
	(c)	Write the ionic equation with state symbols that describes the neutralisation of	

HGV Sec 4E/5NA Science (Chemistry) Prelim Paper 3 2021

any alkali by any acid.

[2]

END OF PAPER Data Sheet

Colours of Some Common Metal Hydroxides

calcium hydroxide	white
copper(II) hydroxide	light-blue
iron(II) hydroxide	green
iron(III) hydroxide	red-brown
lead(II) hydroxide	white
zinc hydroxide	white

The Periodic Table of Elements

The wokur						=			j	fiancium		87	133	carrium	က္သ	35	8	ð	37	36		- d	23	SOCIUM	2 :	-	7 militari	_	΄ ψ			_	
The volume of one mole of any gas is 24 dm ³ at room temperature and pressure (r.t.p.).			actinoids			lanthanoids			1	mulber	20	88	137				88	Ľ	<u>ب</u> ي د	40) N	24	muignesium	<u></u>	3 8	beryllium	Be	4			=	
moke of a							•					89 - 103			lanthanoids		89	~	39	45	SC C	2 2								•			
ıny gas is	I distribution	7	88	139		- 57			ı	Rutherfordium	₹	1	178	hamium	Į	72	91	4	8	48		3 15				reiam		ato	proton				
24 dm³ a	232	₹	8	140	6 6 6	쎯			١	dubnium	8	105	181		.	73	93	Z	4	51	<	: 23				relative apprinc mass	FIBITION	atomic symbol	proton (atomic) number	Key			:
at room te	231	Pa	2	141	Pr	58			ı	Seaborgium	တ္ဆ	106	184	tungaten	€:	74	96	Mo	₽	52	ב	2				mass		<u>8</u>	umber				
mperatur	238	_	න	144	No	0.9			ı	bohrium	9	701	186	menum	ਨਾਂ :	75	- TOTAL CHARGE	TC	43	55	MA	۲										•	
e and pre	nepainium –	S	93	1	Pm Pm	61			1	hassium	댫	108	190	CSATILLED		\neg				83		26								hydrogen 1	Ι -		
ssure (r.t	Pillionium	وح	2	150	SID	29			ı	mailneours	≚	109	192	indium:	≓ `	η	103	골	45	59	Ş	27											Gn
.p.).	americam.	A A	95	152		ස			ı	darmstadtium	<u>ک</u>	110	195	platinum	Ţ	78	106	2	46	59 64	Z	8	-										Group
	curium	ဌ	96	157	G G G	64			ı	muknaganaca	2 0	11	197	0 1	₽ ;	79	108 108	8	47	ğ 40	5	29	•										
	berkeilum	무	97	58	7	ස			1	copernicium	ջ <u>։</u>	112	201							8.53			•										
	oakiomium	Ω	88	163	Dy	66						I		thalkion ;	.	82		5	49	70	G	<u> </u>	27	eluminium	≥ Շ	3 =	boron	œ	5			NI NI	
	n einsteinum sei	Ľ	88	165		67			1	flerovium	IJ	114	207	8 (구 :	82	116	ရှ	25	73	Ge	<u>ئ</u> د	28	slicon	<u> </u>	Z	Calibon	ဂ	6			7	
		. ⊒	8	167	<u>‡</u> П	68	C.	m	ıi	4	ris I	\neg				\dashv				75			-			1		z	7			٧	
	mendeleviu	<u> </u>	101	169		66		. 1	1	livermonu	<u>ر</u>					~				79 -			٢					0	œ			≤	

HG

<u>Sec 4E/5N Science Chemistry Preliminary Exam 2021</u> <u>Answer Scheme</u>

Paper 1 - MCQ Answers

	21.	С	22.	A	23.	С	24.	С	25.	С	26.	В	27.	C	28.	C	29.	В	30.	Α
Γ	31.	D	32.	С	33.	В	34.	D	35.	С	36.	С	37.	В	38.	D	39.	D	40.	В

Paper 3 - Section A - Structured Questions

		Answers	Mark	Remarks for markers
1	a)	A	1M	
	b)	D	1M	
	c)	E	1M	
	d)	В	1M	
	e)	A, B or C	1M	
2	a) _	S C S	2M	Formula for CS ₂ must be correct first to award any mark 1M for bonding electrons between C and S 1M for other electrons of S
	b)	True Falselow boiling point very soluble in watergood electrical conductivity when molten crystalline solid at room	2M	4 correct – 2M 2-3 correct – 1M 1 correct – 0M
	c)	i) 2+ 2-	2M	1M for each ion correctly drawn inclusive of charges 2M total If inner electrons are drawn,
		Mg S S		it must be correct, otherwise no mark is awarded Best practice: 2 crosses for sulfur (or vice versa) should be denoted that it came from magnesium (but marks can be awarded if it's all drawn using the same symbol)

oppository stranger to the	1M 1M 1M 1M 1M 1M 1M	[idea of solid formed] [idea of blocking/covering cross] Words in brackets not necessary as it's mentioned in the question. Temperature or temperature of surroundings. [Accepted]
Thus it requires a lot of energy to overcome them. 3 a) Sulfur is formed and is an insoluble solid. Thus as more sulfur is formed, it blocks the cross from being seen/obscures the cross (OWTTE) b) i) It (Rate of reaction) decreases (as concentration decreases) ii) lower concentration → less reactant particles per unit volume/in a given volume lower frequency of effective collisions iii) any 2 temperature of acid/thiosulfate solution volume of acid volume of thiosulfate solution concentration of acid iv) 1.1 1.0 0.9 0.8 0.7 0.6 0.5 0.4 0.3 0.2 ***	1M 1M 1M 1M	[idea of blocking/covering cross] Words in brackets not necessary as it's mentioned in the question. Temperature or temperature of surroundings. [Accepted]
a) Sulfur is formed and is an insoluble solid. Thus as more sulfur is formed, it blocks the cross from being seen/obscures the cross (OWTTE) b) i) It (Rate of reaction) decreases (as concentration decreases) ii) lower concentration → less reactant particles per unit volume/in a given volume lower frequency of effective collisions iii) any 2 temperature of acid/thiosulfate solution volume of acid volume of thiosulfate solution concentration of acid iv) 1.1 1.0 0.9 0.8 0.7 0.6 0.5 0.4 0.3 0.2 *** ** Sulfur is formed and is an insoluble solid. Thus as more sulfur is formed, it blocks the cross from being seen/obscures the cross (OWTTE) ii) lower concentration → less reactant particles per unit volume for acid solution solution.	1M 1M 1M	[idea of blocking/covering cross] Words in brackets not necessary as it's mentioned in the question. Temperature or temperature of surroundings. [Accepted]
Thus as more sulfur is formed, it blocks the cross from being seen/obscures the cross (OWTTE) b) i) It (Rate of reaction) decreases (as concentration decreases) ii) lower concentration → less reactant particles per unit volume/in a given volume lower frequency of effective collisions iii) any 2 temperature of acid/thiosulfate solution volume of acid volume of thiosulfate solution concentration of acid iv) 1.1 1.0 0.9 0.8 0.7 0.6 0.5 0.4 0.3 0.2	1M 1M 1M	[idea of blocking/covering cross] Words in brackets not necessary as it's mentioned in the question. Temperature or temperature of surroundings. [Accepted]
from being seen/obscures the cross (OWTTE) b) i) It (Rate of reaction) decreases (as concentration decreases) ii) lower concentration → less reactant particles per unit volume/in a given volume lower frequency of effective collisions iii) any 2 temperature of acid/thiosulfate solution volume of acid volume of thiosulfate solution concentration of acid iv) 1v) 1.1 1.0 0.9 0.8 0.7 0.6 0.5 0.4 0.3 0.2	1M 1M	Words in brackets not necessary as it's mentioned in the question. Temperature or temperature of surroundings. [Accepted]
decreases) ii) lower concentration → less reactant particles per unit volume/in a given volume lower frequency of effective collisions iii) any 2 temperature of acid/thiosulfate solution volume of acid volume of thiosulfate solution concentration of acid iv) 1.1 1.0 0.9 0.8 0.7 0.6 0.5 0.4 0.3 0.2	1M 1M	necessary as it's mentioned in the question. Temperature or temperature of surroundings. [Accepted]
unit volume/in a given volume lower frequency of effective collisions iii) any 2 temperature of acid/thiosulfate solution volume of acid volume of thiosulfate solution concentration of acid iv) 1.1 1.0 0.9 0.8 0.7 0.6 0.5 0.4 0.3 0.2	1M	temperature of surroundings. [Accepted]
lower frequency of effective collisions iii) any 2 temperature of acid/thiosulfate solution volume of acid volume of thiosulfate solution concentration of acid iv) 1.1 1.0 0.9 0.8 0.7 0.6 0.5 0.4 0.3 0.2		temperature of surroundings. [Accepted]
iii) any 2 temperature of acid/thiosulfate solution volume of acid volume of thiosulfate solution concentration of acid iv) 1.1 1.0 0.9 0.8 0.7 0.6 0.5 0.4 0.3 0.2	1M	temperature of surroundings. [Accepted]
volume of acid volume of thiosulfate solution concentration of acid iv) 1.1 1.0 0.9 0.8 0.7 0.6 0.5 0.4 0.3 0.2		surroundings. [Accepted]
volume of thiosulfate solution concentration of acid iv) 1.1 1.0 0.9 0.8 0.6 0.5 0.4 0.3 0.2		
iv) 1.1 1.0 0.9 0.8 0.7 0.6 0.5 0.4 0.3 0.2		2M
1.1 1.0 0.9 0.8 0.7 0.6 0.5 0.4 0.3 0.2		214
1.0		
1.0		1M for plots 1M for curve drawn (smooth
0.9 0.8 0.7 0.6 0.5 0.4 0.3 0.2		curve)
0.8 0.7 0.6 0.5 0.4 0.3 0.2		
0.7 0.6 0.5 0.4 0.3 0.2		
0.6 0.5 0.4 0.3 0.2		
0.5 0.4 0.3 0.2		
0.4		
0.3		
0.2		
0.0 0	45	
0 3 10 13 20 23 30 33 40	43	
v) 0.55 to 0.6 mol/dm ³	i	Any acceptable value based on a curve drawn.
	1M	
4 a) +4 - oxidised / increased +2 - reduced / decreased	1M	1M each for oxidation states

	b)	 Forms acid rain → any 1 harmful effect of acid rain leaches nutrients from the soil and harm plant growth acidifies lakes (closed water bodies), streams, rivers and harm aquatic life corrodes buildings and structures made of limestone and metals Or irritates the eyes/ cause breathing difficulties – irritate the lungs/throat / inflammation of the lungs/causes bronchities binds irreversibly/reacts with red blood 	1M	
	:	cells(haemoglobin) and deprive them of oxygen can result if fatigue, headaches, feeling faint or even death		
	c)	i) 114	1M	
		ii) 1.16	1M	(ecf from part(i)-accepted)
		iii) 14.5	1M	(ecf from part(ii)-accepted)
			4 5 #	
5	(a)	i) barium sulfate / BaSO₄	1M	
	<u> </u>	ii) carbon dioxide / CO ₂	1M	4
		iii) magnesium / Mg	1M	4
	ļ	iv) magnesium sulfate MgSO ₄	1M	4
		v) magnesium oxide / MgO OR magnesium hydroxide / Mg(OH) ₂	1M	
5	b)	Ba ²⁺ (aq) + SO ₄ ²⁻ (aq) → BaSO ₄ (s) OR Mg(s) + 2H ⁺ (aq) → Mg ²⁺ (aq) + H ₂ (g) MgCO ₃ (s) + 2H ⁺ (aq) → Mg ²⁺ (aq) + CO ₂ (g) + H ₂ O(<i>l</i>) MgO(s) + 2H ⁺ (aq) → Mg ²⁺ (aq) + H ₂ O(<i>l</i>) Mg(OH) ₂ (s) + 2H ⁺ (aq) → Mg ²⁺ (aq) + 2H ₂ O(<i>l</i>)	1M	MgO and Mg(OH) ₂ are taken as insoluble (due to their low solubility)
	a)	substances element compound mixture F G H I	5M	1M for each tick
	b)	 A mixture has a variable composition of its constituent substances but a compound has a fixed composition of its constituent elements. A mixture has a variable ratio of its components but the elements in a compound are found in a fixed ratio. 		Keywords in bold are necessary to illustrate the concept Same point.

Section B - Free - Response Questions

		Answers	Mark allocation	Remarks
7	a)	Sodium has an electronic configuration of 2,8,1.	1M	
	'	It has 1 valence electron and is in Group I.	1M	
		It has 3 electron shells and is thus in period 3	1M	
	b)	Both of them react with water to form a metal	1M	
		hydroxide (or an alkali) and hydrogen gas.		
		Both of them react with oxygen to form metal		
		oxides.	1M	
		Both of them react with halogens to form metal		
		halides.		Eqn can be done with Na
	Ì			also
		$2K(s) + 2H_2O(l) \rightarrow 2KOH(aq) + H_2(g) OR$	2M	1M for correct and
	ļ	$4K(s) + O_2(g) \rightarrow 2K_2O(s)$	1	balanced equation
		$2K(s) + Cl_2(g) \rightarrow 2KCl(s)$		1M for state symbols
	c)	Sodium is from the group called Alkali metals.	1M	For both. Group I and 0
	-,	Neon is from the group called Noble gases.		not accepted.
		sodium atom is very reactive as its atom has 1		
		valence electron and it loses it easily *fto obtain a	1M	The idea of
		completely filled valence shell.]		reactivity/unreactive
		Neon has a complete (or completely filled) valence	1M	linked to
	ļ	shell and thus is unreactive.		incomplete/complete
				valence shells must be
		*does not need to be mention if a comparison to		stated
		neon is done (with neon having a complete valence		
		shell – linked to it being unreactive)		
8	a)	Isotopes	1M	
	b)	They have different masses as the atoms have		
		different number of neutrons.	1M	
		They have the same chemical properties as they		
		have the same number of valence electrons. (or 7	1M	
		valence electrons.		

	c)	i) Relative atomic mass is the <u>average mass of an atom of the element</u> as compared to 1/12 th the	1M	
		mass of a carbon-12 atom. ii) $\frac{75}{100}$ x 35 + $\frac{25}{100}$ x 37 = 35.5	1M	
	d)	7 100 100	3M	1M for no. of protons 1M for no. of neutrons 1M for correct no. of shells and electrons (nucleus outline drawn or not doesn't matter)
	e)	Mr of $Cl_2 = 71$ No. of moles of $Cl_2 = 142/71$ = 2 Mole ratio > $Cl_2 : COCl_2$ 1 : 1 2 : 2 Volume of phosgene = 2 x 24 = 48 dm ³	1M	
9	a)	 Pipette 25.0cm³ of sodium hydroxide solution into a conical flask. Add a few drops of methyl orange indicator. Titrate the sodium hydroxide solution with a solution of sulfuric acid from a burette. Stop the titration when the indicator change colour. Record the volume of acid used. Repeat the titration using the volume of acid recorded without the indicator. 	1M 1M 1M 1M 1M	A suitable indicator. Universal indicator – Rej. Acid must be correct in order to award this mark If student reverse the substances in the conical flask/burette. It is also accepted.
	b)	No. of moles of sodium hydroxide = 0.250 x 2 = 0.5	1M	
		Mass of NaOH = 0.5 x 40 = 20 g	1M	
	c)	H ⁺ (aq) + OH ⁻ (aq) → H ₂ O(<i>l</i>)	2M	1M for correct balanced eqn. 1M for state symbols